



NOAA Technical Report NMFS SSRF- 742

WATER STRUCTURE AT OCEAN WEATHER STATION V NORTHWESTERN PACIFIC OCEAN, 1966-71

D. M. Husby and G. R. Seckel

October 1980

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service

NOAA TECHNICAL REPORTS

National Marine Fisheries Service, Special Scientific Report—Fisheries

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The Special Scientific Report—Fisheries series was established in 1949. The series carries reports on scientific investigations that document long-term continuing programs of NMFS, or intensive scientific reports on studies of restricted scope. The reports may deal with applied fishery problems. The series is also used as a medium for the publication of bibliographies of a specialized scientific nature.

NOAA Technical Reports NMFS SSRF are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained (unless otherwise noted) from D822, User Services Branch, Environmental Science Information Center, NOAA, Rockville, MD 20852. Recent SSRF's are:

700. Expendable bathythermograph observations from the NMFS/MARAD Ship of Opportunity Program for 1973. By Steven K. Cook. June 1976, iii + 13 p., 10 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; Stock No. 003-017-00382-5.
701. Seasonal abundance and distribution of zooplankton, fish eggs, and fish larvae in the eastern Gulf of Mexico, 1972-74. By Edward D. Houde and Nicholas Chitty. August 1976, iii + 18 p., 14 figs., 5 tables.
702. Length composition of yellowfin, skipjack, and bigeye tunas caught in the eastern tropical Atlantic by American purse seiners. By Gary T. Sakagawa, Athlo L. Coan, and Eugene P. Holzapfel. August 1976, iv + 22 p., 7 figs., 7 tables, 15 app. tables.
703. Aquacultural economics bibliography. By John Vondruska. October 1976, 123 p.
704. The macrofauna of the surf zone off Folly Beach, South Carolina. By William D. Anderson, Jr., James K. Dias, Robert K. Dias, David M. Cupka, and Norman A. Chamberlain. January 1977, iv + 23 p., 2 figs., 31 tables.
705. Migration and dispersion of tagged American lobsters, *Homarus americanus*, on the southern New England continental shelf. By Joseph R. Uzmann, Richard A. Cooper, and Kenneth J. Pecci. January 1977, v + 92 p., 45 figs., 2 tables, 29 app. tables.
706. Food of western North Atlantic tunas (*Thunnus*) and lancetfishes (*Alepisaurus*). By Frances D. Matthews, David M. Damkaer, Leslie W. Knapp, and Bruce B. Collette. January 1977, iii + 19 p., 4 figs., 1 table, 11 app. tables.
707. Monthly temperature and salinity measurements of continental shelf waters of the northwestern Gulf of Mexico, 1963-65. By Robert F. Temple, David L. Harrington, and John A. Martin. February 1977, iii + 26 p., 5 figs., 10 tables.
708. Catch and catch rates of fishes caught by anglers in the St. Andrew Bay system, Florida, and adjacent coastal waters, 1973. By Doyle F. Sutherland. March 1977, iii + 9 p., 2 figs., 9 tables.
709. Expendable bathythermograph observations from the NMFS/MARAD Ship of Opportunity Program for 1974. By Steven K. Cook and Keith A. Hausknecht. April 1977, iv + 45 p., 10 figs., 9 tables, 35 app. figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; Stock No. 003-017-00397-3.
710. Midwater invertebrates from the southeastern Chukchi Sea: Species and abundance in catches incidental to midwater trawling survey of fishes, September-October 1970. By Bruce L. Wing and Nancy Barr. April 1977, iii + 43 p., 1 fig., 2 tables, 2 app. tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; Stock No. 003-020-00130-1.
711. A list of the marine mammals of the world. By Dale W. Rice. April 1977, iii + 15 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; Stock No. 003-020-00134-3.
712. Annual physical and chemical oceanographic cycles of Auke Bay, southeastern Alaska. By Herbert E. Bruce, Douglas R. McLain, and Bruce L. Wing. May 1977, iii + 11 p., 16 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; Stock No. 003-020-00134-3.
713. Current patterns and distribution of river waters in inner Bristol Bay, Alaska. By Richard R. Straty. June 1977, iii + 13 p., 16 figs., 1 table.
714. Wind stress and wind stress curl over the California Current. By Craig S. Nelson. August 1977, iii + 87 p., 18 figs., 1 table, 3 app. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; Stock No. 003-020-00139-4.
715. Bottom obstructions in the southwestern North Atlantic, Gulf of Mexico, and Caribbean Sea. By G. Michael Russell, Abraham J. Barrett, L. Steve Sarbeck, and John H. Wordlaw. September 1977, iii + 21 p., 1 fig., 1 app. table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; Stock No. 003-020-00140-8.
716. Fishes and associated environmental data collected in New York Bight, June 1974-June 1975. By Stuart J. Wilk, Wallace W. Morse, Daniel E. Ralph, and Thomas R. Azarovitz. September 1977, iii + 53 p., 3 figs., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; Stock No. 003-017-0040-4.
717. Gulf of Maine-Georges Bank ichthyoplankton collected on ICNAF larval herring surveys September 1971-February 1975. By John B. Colton, Jr. and Ruth R. Byron. November 1977, iii + 35 p., 9 figs., 14 tables.
718. Surface currents as determined by drift card releases over the continental shelf off central and southern California. By James L. Squire, Jr. December 1977, iii + 12 p., 2 figs.
719. Seasonal description of winds and surface and bottom salinities and temperatures in the northern Gulf of Mexico, October 1972 to January 1976. By Perry A. Thompson, Jr. and Thomas D. Leming. February 1978, iv + 44 p., 43 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; Stock No. 003-017-00414-7.
720. Sea surface temperature distributions obtained off San Diego, California, using an airborne infrared radiometer. By James L. Squire, Jr. March 1978, iii + 30 p., 15 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; Stock No. 003-017-00415-5.
721. National Marine Fisheries Service survey of trace elements in the fishery resource. By R. A. Hall, E. G. Zook, and G. M. Meahurn. March 1978, iii + 313 p., 5 tables, 3 app. figs., 1 app. table.



NOAA Technical Report NMFS SSRF- 742

**WATER STRUCTURE AT
OCEAN WEATHER STATION V,
NORTHWESTERN PACIFIC OCEAN,
1966-71**

D. M. Husby and G. R. Seckel

October 1980

U.S. DEPARTMENT OF COMMERCE

Philip M. Klutznick, Secretary

National Oceanic and Atmospheric Administration

Richard A. Frank, Administrator

National Marine Fisheries Service

Terry L. Leitzell, Assistant Administrator for Fisheries

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

Introduction	1
Processing procedures and analysis	1
The data	1
The isentropic format	2
Quality control of isentropic data	2
Harmonic analysis of isentropic data	2
Statistical characteristics of the harmonic analyses	5
Sampling gaps	12
Results of the harmonic analyses	13
Oceanographic conditions at OWS-V	13
The oceanographic setting	13
Variability and structure 1966-71	14
The water structure and air-sea interaction	24
Summary and conclusions	26
Acknowledgments	29
Literature cited	29
Appendix I	31
Appendix II	50

Figures

1. Sampling record of oceanographic stations at Ocean Weather Station V, 1966-72	3
2. Positions of oceanographic stations taken by U.S. Coast Guard Cutter <i>Chautauqua</i> at Ocean Weather Station V, 4-22 January 1966	4
3. Original and expected salinity and depth values at σ_t 26 and σ_t 27 for the year 1967	6-7
4. Original and expected salinity and depth values at σ_t 26 and σ_t 27 for the year 1971	8-9
5. Standard errors of estimate for harmonic fit of the 1971 temperatures at σ_t 25 to σ_t 27	10
6. Standard errors of estimate for harmonic fit of the 1971 salinities at σ_t 25 to σ_t 27	10
7. Standard errors of estimate for harmonic fit of the 1971 depths at σ_t 25 to σ_t 27	11
8. Temperature-salinity diagrams for day 244 (1 September 1969) based on original isentropic data with temperature-salinity values from the harmonic fit	12
9. Temperature-salinity diagrams for day 182 (1 July 1971) based on original isentropic data with temperature-salinity values from the harmonic fit	12
10. Dynamic topography of the sea surface relative to the 1,000 decibar surface derived from selected stations during the summer of 1957	14
11. Meridional sections, lat. 21° to 42°N, long. 168°E, of salinity and depth versus σ_t based on 67 STD stations occupied 3-11 April 1971 and dynamic topography of sea surface relative to the 1,000 decibar surface along same section	15
12. Sections of salinity versus thermocline anomaly and thermocline anomaly versus depth along right-hand edge of Kuroshio	16
13. Monthly mean surface temperature-salinity relationships at Ocean Weather Station V, 1966-71	17
14. Vertical profiles of expected values of salinity and depth from harmonic series at the first day of each month at Ocean Weather Station V, 1966-71	18-23
15. Absolute magnitude of the harmonic function for the 6-yr series of the depth of σ_t 26.0	24
16. Heat content of the water column from the surface to σ_t 26.0 at Ocean Weather Station V at 30-day intervals, heat content based on change in depth per month at the σ_t 26.0 level, and the depth of the σ_t 26.0 level computed at midmonth from the harmonic series	25
17. Heat content change per month based on the change in mean temperature per month in layer from sea surface to σ_t 26.0 and heat content change per month contributed by net heat exchange across sea surface	27
18. Change per month of the mean temperature from the sea surface to σ_t 26.0 at Ocean Weather Station V computed from the harmonic series	28

Tables

1. Manner of obtaining first and last values in each of the annual harmonic analyses, 1966-71	4
2. Standard errors of estimate for Fourier series summations of temperature, salinity, and depth on σ_t 25.0-27.4 at Ocean Weather Station V, 1966-71	5

3. Heat budget estimates at Ocean Weather Station V for the 6-mo cooling portion of the year, 1 October to 1 April	24
4. Estimates of mean evaporation rates, mean precipitation rates, and evaporation minus precipitation at Ocean Weather Station V	26

Water Structure at Ocean Weather Station V, Northwestern Pacific Ocean, 1966-71

D. M. HUSBY and G. R. SECKEL¹

ABSTRACT

The oceanographic station data obtained at Ocean Weather Station V from 1966 to 1972 by the U.S. Coast Guard have been analyzed and are presented in a form suitable for water structure studies. Temperatures, salinities, and depths are given as a function of density (σ_t). We used harmonic analysis as a curve-fitting technique, to obtain parameters for these properties as a function of time. The harmonic coefficients and interpolated values at the first of each month for the 6-year series are tabulated in an appendix.

We describe the temporal distributions of salinity and depth in terms of the oceanographic setting. At depths greater than σ_t 26, temperature-salinity relationships remain relatively constant in time. Depth variations at these levels are attributed primarily to meanders of the Kuroshio Extension. The surface divergence, as reflected by changes in the depth of σ_t 26, has no annual periodicity. The 6-year record shows that large baroclinic variability occurs at time scales of more than 35 days with largest variability occurring at the interannual scale.

Heat budget estimates show that the effects of local ocean-atmosphere exchange processes are obscured by advected properties. For example, the heat content of the layer above σ_t 26 is primarily determined by the divergence of this layer and anomalies in the mean temperature are produced by heat advection rather than heat exchange across the sea surface.

INTRODUCTION

Monitoring and predicting ocean variability in the fishing areas of the eastern North Pacific are important problems facing the fishery oceanographer. In contrast to the global monitoring of the atmosphere at a network of stations every 6 h, there is no network of monitoring stations in the oceans. Oceanographers are therefore attempting to infer ocean variability from atmospheric forcing, i.e., the exchange of momentum, moisture, and heat between the atmosphere and ocean, that can be calculated from the regularly observed meteorological properties. The temperature structure in the upper layers of the ocean, for example, is affected by heat exchange and wind stress as well as by the changing current field and diffusion. Inferring ocean variability from atmospheric forcing, therefore, is not a trivial problem.

Conditioning of the water reaching the eastern North Pacific Ocean begins in the western North Pacific, an area characterized by a net annual heat loss across the sea surface (Husby and Seckel 1975). Thus, studies of the effects of heat exchange across the sea surface and wind stress on ocean properties and structure must be undertaken upstream of the fishing areas if predictions of anomalous water conditions are to be made. The only time series of concurrent meteorological and oceanographic data for use in such studies are those that were obtained at ocean weather stations. One of these, Ocean Weather Station V (OWS-V) was located at lat. 34°N,

long. 164°E within the net annual heat loss area that is of concern to us.

At OWS-V meteorological observations were obtained from 1951 to 1972 and oceanographic station data from 1966 to 1972. The large-scale air-sea interaction processes derived from the station's meteorological data were described by Husby and Seckel (1975). Our work is a companion paper in which the oceanographic station data are presented in a form suitable for studies of the water structure. Temperatures, salinities, and depths at selected density (σ_t) levels are analyzed and presented as a function of time. Harmonic analysis is used as a curve-fitting technique, to obtain parameters for these properties as a function of time. Results of the analyses are presented in Appendices I and II. Finally, the time variations in the distribution of properties, the water structure, are described in terms of the oceanographic setting at OWS-V.

PROCESSING PROCEDURES AND ANALYSIS

The Data

Oceanographic sampling at OWS-V began in January 1966 with daily Nansen bottle casts by the U.S. Coast Guard on alternate 3-wk patrols. In March 1968 the program was expanded to daily observations on each patrol. The temperature and salinity were sampled at the sea surface and at the depths of 10, 30, 50, 75, 100, 150, 200, 300, 400, 600, 800, 1,000, and 1,500 m. The observations and initial data processing methods were described by Husby (1968).

¹Pacific Environmental Group, National Marine Fisheries Service, NOAA, c/o Fleet Numerical Oceanography Center, Monterey, CA 93940.

From 1966 to March 1968 the sampling was intermittent (Fig. 1). Subsequently, until January 1972, daily sampling became nearly continuous with isolated gaps of up to 20 days. In addition to the temporal gaps, stations were occasionally occupied outside the nominal location, a 10 nautical mile (nmi) square centered on lat. 34°N, long. 164°E (Fig. 2). We, however, used all observations within a 60 nmi square centered on this location. Sixty-seven percent of the stations were occupied within the 10 nmi square and 95% within the 60 nmi square.

The oceanographic station data were obtained from the National Oceanographic Data Center (NODC) of the National Oceanic and Atmospheric Administration in two different formats: 1) temperature and salinity at observed and standard (interpolated) depths, and 2) temperature, salinity, and depth at increments of 0.2 sigma- t (σ_t) units between the surface and the deepest observation (isentropic format). A cubic spline interpolation function was used to obtain the values of temperature, salinity, and depth at the desired σ_t values.²

The Isentropic Format

The study of changes in water properties on surfaces of constant potential density or σ^θ , called isentropic analysis, was introduced to oceanography by Parr (1938) and Montgomery (1938) and has been a valuable tool in descriptive oceanography. The assumptions underlying isentropic analysis are that mixing or interchange of water masses on a constant density surface proceed with a minimum of change in the potential energy and entropy of the system and that surfaces of constant density are the preferred surfaces along which mixing occurs. The method has been used in the identification and tracing of water masses and also lends itself to the analysis of temporal changes in the water structure as was done, for example, at OWS-P by Tabata (1965). In order to pursue the latter aspect of isentropic analysis the temperature, salinity, and depth are presented as a function of σ_t in this report.

Quality Control of Isentropic Data

The initial step of quality control is concerned with the interpolated temperatures and salinities at designated σ_t values for each station received from NODC. The spline interpolation function calculates the temperature and salinity at designated σ_t values from the two relationships: 1) observed temperature versus σ_t and 2) salinity versus σ_t with σ_t as the independent variable. A σ_t value was computed from the interpolated temperature and salinity at each designated σ_t using the relationships (1) and (2). When the calculated σ_t differed from the designated by $> \pm 0.02 \sigma_t$ units at more than one level, both interpolated and observed data were used to plot two temperature-salinity (T-S) dia-

grams for the station. The T-S curve drawn through the observed values together with the temperature-depth curve was used to correct the interpolated values obtained from the spline interpolation or to determine whether the station should be rejected. These quality control procedures frequently had to be used for values that were erroneously produced by the spline interpolation because of an inadequate sampling interval in the vicinity of the salinity minimum (σ_t 26.8). Out of a total of 1,067 stations, 496 stations were corrected during this quality control procedure.

Harmonic Analysis of Isentropic Data

Harmonic analysis was used as a curve-fitting technique to summarize the daily oceanographic data in a manner that lends itself to studies of temporal changes in the water structure. The resulting analytical expressions yield smoothed estimates of the water properties for any time of the analysis period.

Harmonic (Fourier) analysis was performed on the isentropic data for each year from 1966 through 1971 with a fundamental period of 366 days beginning on 1 January of one year and ending on 1 January of the next year. In these data sets, values at the beginning and end of the annual series were usually not equal and to facilitate the curve-fitting procedure the harmonic analyses were performed on the residuals produced by subtracting the linear trend (day 1 to day 366) from the observed value:

$$R(t) = f(t) - [f(1) + M \cdot (t - 1)] \quad (1)$$

where $f(t)$ is the value of the property (temperature, salinity, or depth) at a constant σ_t at day t , and $t = 1$ to 366. The slope of the straight line fitted to the values at day 1 and day 366 is

$$M = [f(366) - f(1)]/365. \quad (2)$$

The expression for the fitted curve then becomes

$$E(t) = f(1) + M \cdot (t - 1) + F_k(t) \quad (3)$$

$$\text{where } F_k(t) = A_0 + \sum_n \left(A_n \cos \frac{2n\pi t}{T} + B_n \sin \frac{2n\pi t}{T} \right), \quad (4)$$

$$n = 1, 2, \dots, k$$

is the Fourier series derived from the residuals, $R(t)$. The coefficients A_0 , A_n , and B_n were evaluated by use of a standard computer program.

Data gaps in the time series were filled by linear interpolation between observed values in order to satisfy the program requirement of equal time intervals between data. When no observations were available for the first or 366th day of the year, data from the first (last) actual station of the year was extrapolated backward (forward) to the missing day. For the 1969 and 1971 analyses, the values for the first day were computed from a harmonic

²Hamilton, D. 1973. Isentropic analysis and spline interpolation. Natl. Oceanogr. Data Cent. Tech. Rep., 29 p.




Figure 1.—Sampling record of daily oceanographic stations at Ocean Weather Station V, 1966-71.




Figure 2.—Positions of oceanographic stations taken by U.S. Coast Guard Cutter *Chautauqua* at Ocean Weather Station V, 4-22 January 1966. Dashed line encloses a 10-nmi square and solid line encloses a 60-nmi square around Ocean Weather Station V.

analysis of the 366 day period beginning and ending on 2 July. There were sufficient data from July 1968 to July 1971 to use the expected values from these analyses as the interpolated values for the first days of 1969 and 1971 (Fig. 1). The values used for the 1st and 366th days of the annual analyses at each level are listed in Appendix I. Table 1 shows the manner in which these values were obtained.

Before proceeding with the harmonic analyses of the complete data set limited analyses were made as a quality control procedure. Harmonic coefficients were derived to the 12th harmonic for the salinity and depth series at the σ_t 26 and 27 levels of each year. Equation (3) was then used to evaluate and plot the fitted curve

Table 1.—Manner of obtaining first and last values in each of the annual harmonic analyses, 1966-71.

Year	Day = 1	Day = 366
1966	extrapolation from 4 Feb. 1966	observation on 1 Jan. 1967
1967	observation on 1 Jan. 1967	extrapolation from 24 Nov. 1967
1968	extrapolation from 1 Mar. 1968	harmonic analysis from 2 July 1968 to 2 July 1969
1969	harmonic analysis from 2 July 1968 to 2 July 1969	observation on 1 Jan. 1970
1970	observation on 1 Jan. 1970	harmonic analysis from 2 July 1970 to 2 July 1971
1971	harmonic analysis from 2 July 1970 to 2 July 1971	observation on 1 Jan. 1972

together with the isentropic data. Examples of these curves for 1967 (a year with large data gaps) and 1971 (a year with small data gaps) are shown in Figures 3 and 4.

Questionable values are revealed as large discrepancies from the fitted curve. The questionable values were usually associated with the previously mentioned poor vertical sampling interval at individual stations or with stations that were occupied outside the 60 nmi quadrangle about the nominal position of OWS-V (Fig. 2). Using this quality control procedure, 38 stations were rejected from the 6-yr series.

After this quality control procedure the harmonic coefficients were calculated for the temperature, salinity, and depth at σ_t levels between 22.0 and 27.4 with intervals of 0.2 σ_t units, as well as for the surface temperature, salinity, and σ_t . Although the analyses were carried out to the 52d harmonic, resolving fluctuations with a 7-day duration, only the first 12 harmonic coefficients are listed in Appendix I. Also listed are the slope M , and initial values, $f(1)$, for the linear trend at each level. Thus, using the values of Appendix I in Equation (3), expected values for any day of the year for any of the given levels can be obtained.

Because of the seasonal variation in the density of the upper layers, properties at the lower σ_t values will not occur throughout the year. For these situations the dates of the first and last observation are given in the tabulation of Appendix I and derived values will be valid only for the duration bounded by these dates.

Statistical Characteristics of the Harmonic Analyses

Figures 3 and 4 show qualitatively that the fitted curves follow the observed values very well, as they should, because a Fourier series approximation is a least

squares fit regardless of the number of harmonics used in the summation (Jenkins and Watts 1968). A quantitative measure of the goodness of fit is provided by the unexplained variance,

$$S_e^2 = \frac{\sum_{t=1}^T [f(t) - E_k(t)]^2}{T-1} \quad (5)$$


Equation (5) is an estimate of the mean square error or variance of the difference between the observed values and the Fourier series approximation. With an increasing number of harmonics used in the approximation, S_e , the standard error of the estimate, should decrease. These quantitative aspects are presented in Table 2 and illustrated in Figures 5, 6, and 7 by graphs of S_e as a function of the number of harmonics used in the analyses of the temperature, salinity, and depth for 1971.

Considering the temperature and salinity graphs (Figs. 5, 6) first, there is a marked reduction in S_e at the shallower levels as n increases to 6, S_e then decreases slowly with increasing n . For example, at σ_t 25.2, S_e for the temperature is 0.28°C at $n = 1$, 0.135°C at $n = 6$, and 0.12°C at $n = 20$. In the deeper layers the decrease in S_e with increasing harmonic used in the fitting procedure is relatively small. For example, at σ_t 26.4, S_e for the temperature is 0.13°C at $n = 1$, 0.115°C at $n = 6$ and 0.11°C at $n = 20$. The greater variability of the properties at the shallower levels may reflect, in part, low frequency variability in the air-sea interaction processes directly affecting the upper layer of the ocean to approximately the σ_t 25.8 level.

Figure 7 shows that the standard error of estimate for

Table 2.—Standard errors of estimate, S_e , for Fourier series summations ($n = 12$) on σ_t 25.0-27.4 levels of temperature, °C, (upper panel); salinity, ‰, (middle panel); and depth, m, (lower panel). S_e was computed only for times of actual observations, excluding interpolated values, 1966-71.

Year	Level												
	25.0	25.2	25.4	25.6	25.8	26.0	26.2	26.4	26.6	26.8	27.0	27.2	27.4
1966	0.070	0.079	0.114	0.080	0.080	0.103	0.118	0.129	0.144	0.093	0.067	0.059	0.060
1967	0.079	.082	.079	.059	.058	.058	.068	.085	.109	.089	.044	.037	.027
1968	.135	.124	.122	.086	.097	.091	.118	.151	.151	.116	.069	.069	.050
1969	.144	.131	.142	.119	.100	.128	.129	.132	.172	.157	.105	.067	.064
1970	.150	.115	.125	.099	.060	.059	.075	.102	.097	.093	.078	.047	.033
1971	.096	.123	.141	.144	.111	.081	.072	.110	.133	.128	.063	.087	.075
1966	0.022	0.025	0.034	0.022	0.020	0.026	0.026	0.024	0.026	0.016	0.011	0.008	0.008
1967	.026	.025	.023	.017	.015	.015	.016	.018	.019	.015	.008	.005	.004
1968	.044	.039	.038	.025	.026	.024	.028	.032	.029	.020	.010	.011	.008
1969	.046	.041	.042	.034	.027	.033	.031	.029	.032	.026	.016	.011	.011
1970	.047	.036	.037	.028	.016	.015	.017	.022	.018	.015	.011	.006	.005
1971	.030	.038	.041	.041	.030	.021	.017	.020	.024	.021	.009	.013	.010
1966	5.7	10.9	21.8	24.6	23.2	23.5	24.1	24.1	24.0	24.4	26.0	28.4	35.9
1967	9.1	8.6	13.6	22.7	26.4	24.0	20.9	19.5	18.0	22.1	19.9	17.9	23.0
1968	10.7	12.9	15.5	35.5	38.5	40.7	33.6	28.0	25.4	24.7	27.7	35.8	37.1
1969	8.1	13.8	21.7	29.8	32.6	31.6	25.6	23.5	24.0	21.0	22.2	23.9	22.8
1970	7.9	6.1	10.7	15.2	19.5	22.9	20.1	18.3	17.4	16.4	17.0	16.7	27.8
1971	5.2	11.8	15.6	27.4	29.7	24.8	23.0	22.2	19.2	19.3	19.6	20.7	25.9







Figure 3.—Original salinity values (x--x) (upper panel) and original depth values (x--x) (lower panel) and expected values (y--y) from harmonic series (to 12th harmonic) at $\sigma_t 26$ (upper curves) and at $\sigma_t 27$ (lower curves) for the year 1967.



JULIAN DAY

SALINITY ‰




Figure 4.—Original salinity values (x--x) (upper panel) and original depth values (x--x) (lower panel) and expected values (y--y) from harmonic series (to 12th harmonic) at σ_t 26 (upper curves) and at σ_t 27 (lower curves) for the year 1971.





Figure 5.—Standard errors of estimate for harmonic fit of the 1971 temperatures at σ_t 25 to σ_t 25.8 (left panel) and at σ_t 26 to σ_t 27 (right panel).






Figure 7.—Standard errors of estimate for harmonic fit of the 1971 depths at σ_t 25 to σ_t 25.8 (left panel) and at σ_t 26 to σ_t 27 (right panel).

the depth is relatively large at all levels for low harmonics. At the 26.0, 26.2, and 26.4 σ_t levels significant improvement in the standard error continues up to $n = 10$. Relatively high variance at σ_t 26.6, 26.8 and 27 for $n = 6$ indicates that low frequency baroclinicity, probably caused by eddies and current meanders, penetrates deeper than 800 m. S_e approaches an asymptotic value of about 20 m in the deeper σ_t layers indicating the magnitude of the residual variance associated with short-term variability and sampling error.

One would assume that the variance of temperature and salinity decreases with increasing σ_t . Figures 5 and 6 and the tabulated variances of the temperature and salinity (Table 2) show that this assumption is not correct. There is a tendency for S_e to decline from higher values in the upper layers to lower values at σ_t 25.8 or 26.2, then to rise again to higher values at σ_t 26.4, 26.6, or 26.8 before finally declining to the lowest values at σ_t 27, 27.2, and 27.4. In other words, the T-S relationships are less variable below the level of seasonal influence than in the deeper layers at and above the salinity minimum. The reason for this curious tendency may be the uncertainty in the T-S relationship introduced by the inadequate vertical sampling interval (200 m at σ_t ,

26.4-26.8) that was previously mentioned. It is also possible that the relatively large variance at the σ_t 26.4-26.8 levels reflects active mixing of water masses upstream.

Depths derived from the harmonic curve fitting can produce density inversions in the seasonal pycnocline during spring and summer months. These inversions are small and are due to the uncertainty of the harmonic fit being of the same order of magnitude as the depth increment between σ_t levels from the surface to the σ_t 25.2 level range from 1 to 5 m. The standard errors of estimate listed in Table 2 generally are larger than these values and range from 5 to 14 m. These uncertainties in the depth estimates are unavoidable with the method employed, but the errors produced in computations of the integrated properties of the water column, e.g., mean temperature or heat content, will not be large since the depth increments in the upper layer are so small. The levels producing inversions can be deleted in these computations without significantly affecting the results. For example, consider the mean temperature of the water column from the surface to the σ_t 26.0 level on 2 days at OWS-V representing summer conditions, 1 September 1969 (Fig. 8) and 1 July 1971 (Fig. 9). The expected depth values for 1 September 1969 revealed an inversion at σ_t 22.8 and 23.0. Deleting these two levels, the mean temperature was calculated to be 17.20°C, compared with a value of 17.24°C calculated from the

Figure 6.—Standard errors of estimate for harmonic fit of the 1971 salinities at σ_t 25 to σ_t 25.8 (left panel) and at σ_t 26 to σ_t 27 (right panel).




Figure 8.—Temperature-salinity diagram for day 244 (1 September 1969) based on original isentropic data (o) with temperature-salinity values from the harmonic fit (x). The numerals give the depth in meters from the harmonic fit (x). The numbers in parentheses represent the difference between the depth of the original isentropic values and those from the harmonic fit.

original isentropic data. For 1 July 1971 the expected isentropic values showed no inversions, but the comparison was made by deleting the σ_t 24.0 and σ_t 24.2 levels at 3 and 9 m, respectively, giving a mean temperature of 15.31°C, which was identical with that calculated from the original isentropic data and compared with 15.25°C before the data were deleted.

Sampling Gaps

Figure 1 shows that there were gaps lasting up to several months in the oceanographic data series. From 1966 to the winter of 1968 oceanographic stations at OWS-V were occupied only on alternate Coast Guard patrols. Subsequent to this time there were gaps of up to




Figure 9.—Temperature-salinity diagram for day 182 (1 July 1971) based on original isentropic data (o) with temperature-salinity values from the harmonic fit (x). The numerals give the depth in meters from the original isentropic data. The numbers in parentheses represent the difference between the depth of the original isentropic values and those from the harmonic fit.

3 wk in the series. The data gaps were bridged by linear interpolation, as previously described, solely to facilitate the analysis procedure. Therefore, although Equation (3) will provide interpolated values for the gaps, the significance of such values is limited. The limitation is related to the Nyquist frequency,

$$f = \frac{1}{2\Delta t} = \frac{n}{T} \quad (6)$$

specifying the highest harmonic to which, in theory, Fourier analysis can be carried out when the data interval is Δt (Jenkins and Watts 1968). Thus, with $\Delta t = T/2n$, the longest 1966 data gap of about 40 days permits interpolation if the summation in Equation (4) is carried out to $n = 4$. The longest 1967 gap of about 60 days limits meaningful summation in Equation (4) to $n = 3$.

The gap between day 325 of 1967 and day 55 of 1968 comprised almost 90 days. Again, in order to facilitate the analysis procedure the values for day 60 of 1968 were

extrapolated backwards to day 1. Consequently, expected values for day 1 through day 60 have no validity. Interpolated values could have been obtained if the harmonic analyses were carried out from July 1967 to July 1968. A 90-day gap, however, permits meaningful summation only to $n = 2$ in Equation (3) which, as is evident from Figures 5, 6, and 7, would resolve only a small portion of the variance.

After the winter of 1968, the longest gap in the data series of about 20 days occurred in 1969, limiting significant interpolation for this gap to $n = 9$ in the summation of Equation (4). In general, all coefficients to $n = 12$ that are tabulated in Appendix I can be used to evaluate the harmonic function for time spans, including those in 1966 and 1967, where data gaps are 15 days or less.

Results of the Harmonic Analyses

Within the limitations discussed in the previous section the harmonic coefficients tabulated in Appendix I can be used in Equation (3) to obtain smoothed values of the temperature, salinity, and depth at any of the designated levels from the sea surface to $\sigma_t 27.4$. This calculation was performed and the results were tabulated in Appendix II for the first of each month of each of the years 1966 through 1971. The values for the 1966 and 1967 periods were computed by summing the harmonic series to the 3d harmonic, while the values for the succeeding years were computed to the 12th harmonic. In addition, the values for the 366th day for each analysis period are included to allow the month-to-month changes in properties to be computed for the entire year. Because harmonic analyses were performed for individual years, the expected values on the 366th day of one year may not match up with the corresponding values on the first day of the following year.

In order to facilitate the computation of dynamic heights at the surface and acceleration potentials at σ_t levels relative to 1,000 dB, interpolated temperatures, salinities, and σ_t values at 1,000 m are included in the tabulations of Appendix II.

OCEANOGRAPHIC CONDITIONS AT OWS-V

The Oceanographic Setting

In addition to heat exchange and wind induced flow processes, variability in surface properties and vertical structure is affected by shifts of the ocean system relative to the geographically fixed location, variations in the intensity of currents, and eddies. A description of the oceanographic setting of OWS-V, therefore, will aid qualitative interpretations of the low frequency oceanographic variations in the time series presented in this report.

OWS-V is located in waters downstream of the Kuroshio, called the Kuroshio Extension (Sverdrup et al. 1942). Summaries of the oceanographic conditions of this region have been given by Masuzawa (1972) and

Kawai (1972). These articles supplement the earlier work of Fleming (1955), Muromtsev (1958), Uda (1963), Barkley (1968), and others.

The surface distributions of temperature and salinity of the North Pacific Ocean have been presented in numerous atlases and other publications such as that produced by the Japan Hydrographic Association (1975). These show that isopleths of temperature and salinity are generally zonal in the vicinity of OWS-V. Meridional temperature gradients range from about 0.4°C per degree of latitude in summer to 0.9°C per degree of latitude in winter. Salinities at OWS-V tend to be highest in late winter and early spring when the meridional gradient is about 0.04‰ per degree of latitude. Lowest salinities occur from July to October when the meridional gradient is about 0.09‰ per degree of latitude.

Charts of the dynamic topography of the North Pacific Ocean prepared by Reid (1961) and Wyrski (1975) show that OWS-V is located in the eastward extension of the Kuroshio. These charts, however, are based on smoothed, long-term averages and do not exhibit the nature of the flow near OWS-V that materially affects the variability of oceanographic properties at the station. The dynamic topography presented by Kawai (1972), based on observations during the summer and fall of 1957, is reproduced in Figure 10. The tightly packed isopleths of dynamic height show the meandering nature of the Kuroshio Extension between lat. 32° and 38°N. Eddies abound both to the south and to the north of this current.

The existence of eddylike features associated with the Kuroshio current system has been well documented in the literature. The confluence area of the Kuroshio and Oyashio has been called the "Transition Area" by Uda (1938) or the "Perturbed Area" by Kawai (1972). Barkley (1968) drew attention to the eddy structure of this area, and Bernstein and White (1977) have shown that the baroclinic mesoscale eddy energy in this area is an order of magnitude higher than that east of long. 170°W.

In the illustration (Fig. 10), OWS-V is located within the meandering current. It is conceivable that at various times the current could pass either to the north or to the south of the station, but OWS-V would remain in an area of eddy-induced variability. This variability makes analysis of the vertical temperature and salinity distribution difficult if these properties are presented as a function of depth. However, if these properties are presented as a function of density (isentropic format) current and eddy-induced variability is reflected only in the depth distribution of isopycnals.

Meridional and zonal sections of the salinity and depth as a function of σ_t are useful in describing the baroclinic setting. A meridional section of hydrographic stations was occupied in April 1971 at long. 168°E (Roden 1972). Although this section lies to the east of OWS-V, it reflects the characteristic meridional distributions to be found at the longitude of the station. The dominant feature in the salinity section (Fig. 11, upper




Figure 10.—Dynamic topography (dynamic meters) of the sea surface relative to the 1,000 decibar surface derived from selected stations during the summer of 1957 (from Kawai 1972). The location of Ocean Weather Station V is denoted by an "X".

panel) is the constancy of σ_t of the salinity minimum layer. This layer protrudes southward from the subarctic front between lat. 41° and 42°N and is identified with the North Pacific intermediate water (Sverdrup et al. 1942; Reid 1965). During the time of this section, at the latitude of OWS-V, the salinity decreased monotonically from the surface to the depth of the salinity minimum. Farther south, however, shallow salinity maxima occurred.

The distribution of depth as a function of σ_t in the section (Fig. 11, middle panel) reflects the baroclinicity of the zonal flow and embedded eddies. The bottom panel of Figure 11 shows that the principal zonal geostrophic flow occurs between lat. 29° and 35°N. Given the meandering nature of this current (Fig. 10) it is possible that the vertical density structure at OWS-V may be similar to that found within, to the north, or to the south of the current.

Useful zonal sections have been produced by Masuzawa (1972) from a selection of 18 stations during the summer season on the south side of the Kuroshio and Kuroshio Extension. The distribution of thermosteric anomaly versus depth and salinity versus thermosteric anomaly are reproduced in Figure 12. [Surfaces of thermosteric anomaly are parallel to surfaces of σ_t (Montgomery and Wooster 1954).] The middle portion of this section from long. 130° to 170°E runs along approximately lat. 32°N. Note the differences in the distribution of salinity above σ_t 26.0. East of long. 165°E the highest salinities occur at or near the surface, identifying the North Pacific central water. West of long. 160°E a shallow salinity maximum occurs at about σ_t 25. This subtropical water structure with relatively low




salinity surface water is characteristic of the Kuroshio during summer (Masuzawa 1972). At the longitude of OWS-V, the thermosteric anomaly of 200 c/t (about σ_t 26) is at 500 m (Fig. 12), but at long. 168°E and at the same latitude as OWS-V σ_t 26 is near 200 m (Fig. 11). This difference is consistent with the meanders of the current shown in Figure 10.

Variability and Structure 1966-71

The seasonal variability of the sea surface temperature and salinity for the 6 years of hydrographic observations are shown in the T-S diagrams of Figure 13. Six-year averages provide the reference T-S relationships with temperatures ranging from about 16°C in March to 26°C in August and salinities ranging from about 34.73‰ in April to 34.43‰ in September. Departures from the average pattern during each year appear to be more pronounced in the salinity than in the temperature. For example, from June to December 1966 salinities were up to 0.2‰ higher. Higher salinities also occurred during spring and early summer of 1968. Salinities were lower than the average during much of the summer and fall of 1969 and again from June to August of 1971.

A major departure from the average temperature, lasting several months, occurred in 1969-70. Temperatures were 1°C or more below average from

Figure 11.—Meridional sections, lat. 21° to 42°N, long. 168°E, based on 67 STD stations occupied 3-11 April 1971. Upper panel: Salinity (‰) vs. σ_t values. Middle panel: Depth (m) vs. σ_t values. Bottom panel: Dynamic topography (dyn. m.) at the sea surface relative to 1,000 decibar surface.



DEGREES NORTH LATITUDE

line along southern edge of Kuroshio




Figure 12.—Section of salinity (σ_t) versus thermocline anomaly (δ_t) along right-hand edge of Kuroshio (upper panel). Section of thermocline anomaly versus depth (m) along right-hand edge of Kuroshio (lower panel). (From Masuzawa 1972.)

September 1969 to June 1970. In March 1970 the sea-surface temperature was 2.5°C below the 6 yr average. Above-average temperatures occurred in the winters of 1966 and 1969.

The time variability in the vertical structure at OWS-V as reflected by the smoothed isopleths of salinity and depth versus σ_t is shown in Figure 14. Harmonic functions were evaluated at the first of each month using 12 harmonics for 1968-71 and only the first 3 harmonics for 1966 and 1967. The isopleths of salinity and depth are not shown for the time from 1 November 1967 to 1 March 1968 due to the large data gap during this time. However, the expected surface salinity and density for day 331 of 1967 and days 1 and 31 of 1968 are plotted and were derived from the harmonic coefficients for the surface temperature and salinity at OWS-V (Yong 1971).

The common feature for all years in the isentropic

salinity distribution is the constancy of salinity versus σ_t at depths below the σ_t 26 level reflecting a constancy of T-S relationships. Note the constancy of the salinity minimum layer at σ_t 26.8, a feature that was also apparent in both the meridional and zonal sections shown in Figures 11 and 12.

Time variability in the salinity becomes pronounced above the σ_t 26 level and is associated, in part, with the seasonal change in the surface density. Maximum density at the surface occurs in March or April and minimum density in August or September. In 1970, for example, the maximum surface density was only a little less than σ_t 26. During the summer when the surface salinity is lower than in winter, a subsurface salinity maximum forms. In winter the highest salinity of 34.6 or 34.7‰ occurs at the surface. The structure of the subsurface salinity maximum is variable and may appear as




Figure 13.—Monthly mean surface temperature-salinity relationships at Ocean Weather Station V, 1966-71, connected by dashed lines. Arabic numerals represent months. Solid line connects 6-yr mean values. Roman numerals on the 1966 plot give months for the mean values.

well-developed cells as in June to August 1968, August 1969, and November 1971.

The baroclinicity of the area is reflected by the depth of σ_t levels. Time variability penetrates to the σ_t level. The depth of σ_t 26, the level below the seasonal temperature and salinity variability, ranges from about 200 to 400 m. These depth changes are not seasonal but show longer term variations. For example, during much of 1966 σ_t 26 was deeper than 300 m; during 1970 it remained near 200 m. Superimposed on these variations are perturbations of several months duration as in July to September 1969. Amplitudes of the harmonic series, $C_n = (A_n^2 + B_n^2)^{1/2}$, of the depth of the σ_t 26 level for the entire 6-yr record (Fig. 15) show that the greatest portion of the variance, $C_n > 30$ m; occurs on the interannual time scale. Beyond the 39th harmonic (56 day period) the absolute magnitudes of the amplitudes remain < 10 m. Thus, changes in the depth of the pycnocline, as reflected by the σ_t 26 level, are climate-scale phenomena.

In terms of the oceanographic setting of OWS-V, the depth variations reflect north-south shifts of the Kuroshio Extension and/or passage of baroclinic eddies. The salinity variability above the σ_t 26 level permits qualitative interpretation of the observed depth changes. Whenever the depth of σ_t 26 increases to > 300 m (see dashed lines in salinity sections, Fig. 14) there is a concurrent increase in the salinity of the shallow maximum, and during the winter, an increase in the surface salinity. Increases in depths began in February and June 1966, May 1968, April and July 1969, March and November 1971. A shallow salinity maximum is characteristic of Kuroshio water (Fig. 12). Thus, the coincident increases in salinity in the shallow maximum, or at the surface during the winter with increasing depths of σ_t 26 indicate a northward meander of the Kuroshio Extension.

In 1970, when σ_t 26 shoaled to < 300 m, the salinity in the shallow maximum during spring, summer, and fall was relatively high in contrast to that in 1967 when the





Figure 14.—Vertical profiles of salinity (upper panel) and depth (lower panel), with σ_t as ordinate, at Ocean Weather Station V, 1966-71. Values were calculated for first day of each month from the harmonic series carried out to the 3rd harmonic in 1966 and 1967 and to the 12th harmonic in 1968 through 1971. Uppermost solid line connects surface σ_t values. The contour intervals are 0.1‰ for salinity and 100 m for depth. Dashed curves on salinity plot represent 100 m and 300 m isopleths.





Figure 11.—Continued.





Figure 14.—Continued.





Figure 14.—Continued.





Figure 14.—Continued.





Figure 14.—Continued.




Figure 15.—Absolute magnitude of the amplitudes of the harmonic function for the 6-yr series of the depth of σ_t 26.0. Procedure was identical to that used in the annual analyses.

depth of σ_t 26 was also shallow. The high salinity indicates the presence of Kuroshio water at OWS-V although the shallow depth of σ_t 26 did not indicate a northward shift of the Kuroshio Extension. An explanation may be a weakened baroclinic flow during 1970 or a residual consequence of processes upstream of the station that also caused the lower surface temperatures at OWS-V during the first half of 1970.

The Water Structure and Air-Sea Interaction

A study of the effects of atmospheric forcing on ocean variability at OWS-V is not within the scope of this report. Nevertheless, some inferences can be made about the changes described above in the light of the air-sea interactions previously published (Husby and Seckel 1975). For example, principal heat loss from the sea surface occurs during the 6 mo from 1 October to 1 April, and one can inquire about the change in heat content of the water column between these dates.

Consider the layer above σ_t 26 in which temperature and salinity show seasonal changes. The heat content of this layer is $\rho c_p \bar{\theta} z$ where ρ is the density of the water, c_p is the specific heat at constant pressure, $\bar{\theta}$ is the mean temperature of the layer and z is the depth of the layer. With ρ and c_p assumed constant, the heat content of the layer changes because of changes in the thickness of the layer (divergence) and because of changes in the mean temperature.

$$\Delta H = \rho c_p \Delta(\bar{\theta} z) = \rho c_p [\bar{\theta} \Delta z + z \Delta \bar{\theta}]. \quad (7)$$

Changes in the last term are caused by the heat exchange across the sea surface, advection, and diffusion.

In Table 3, the heat content change, the heat content change due to the mean temperature change, and the total heat exchange across the sea surface are listed for

the fall and winter cooling seasons from 1966 to 1971. Note the large interannual variability in the heat content changes with the layer actually gaining heat during the 1967-68 season when the total heat loss across the sea surface was anomalously high. Not only are these changes much larger than can be accounted for by the total heat exchange, they also bear no resemblance to the year to year changes of this atmospheric forcing process. The heat content change due to the change of mean temperature has the same order of magnitude as the total heat exchange but again, the interannual variability does not resemble that of the atmospheric forcing process. For example, the total heat exchange in the 1968-69 season was less than half that of the previous year, and yet the difference in the heat content change due to change in mean temperature was only 8 kcal cm^{-2} . Evidently other processes such as heat advection, play an important part in the change of mean temperature.

The 6-yr series of heat content of the layer to the σ_t 26 level (Fig. 16) shows that during 1966 the heat content was high, and during 1970 it was relatively low with irregular fluctuations between these years. A seasonal

Table 3.—Heat budget estimates at Ocean Water Station V for the 6-mos cooling portion of the year, 1 October to 1 April.

	A ¹ (kcal cm^{-2})	B ² (kcal cm^{-2})	C ³ (kcal cm^{-2})
1 Oct. 1966-1 Apr. 1967	-228.6	-93.0	-62.3
1 Oct. 1967-1 Apr. 1968	96.1	-76.4	-69.8
1 Oct. 1968-1 Apr. 1969	-137.1	-68.5	-33.1
1 Oct. 1969-1 Apr. 1970	-82.1	-51.7	-48.4
1 Oct. 1970-1 Apr. 1971	-139.6	-46.7	-53.6

¹Heat content change in the layer from surface to σ_t 26.

²Heat content change due to the change in mean temperature in layer from the surface to σ_t 26 (see text).

³Total heat exchange across sea surface.




Figure 16.—Upper panel: Heat content (kcal cm^{-2}) of the water column from surface to $\sigma_t 26.0$ at Ocean Weather Station V at 30-day intervals (solid line), and the heat content (kcal cm^{-2}) based on the change in depth per month of the $\sigma_t 26.0$ level (dashed line) (see text). Lower panel: Depth (m) of the $\sigma_t 26.0$ level computed at midmonth from the harmonic series.

pattern is not discernible. The 6-yr series of the depth of σ_t 26 is shown in the lower panel indicating that the fluctuations in depth of the lower boundary of the layer are coherent with those in heat content: high heat content corresponds to greater thickness of the layer and lower heat content with reduced thickness. The coherence is clearly shown by the dashed line in Figure 16, representing the heat content variability due to changes in thickness ($\bar{\theta}\Delta z$) only. The values were obtained by sequentially adding the monthly changes ($\bar{\theta}\Delta z$) to the initial heat content on day 1, 1966. (The series was reinitialized on day 61, 1968.)

The seasonal aspects of the change of heat content per month due to change in the mean temperature in the layer above σ_t 26.0 ($\bar{z}\Delta\theta$), and the monthly heat exchange across the sea surface are shown in Figure 17. The two quantities plotted are of equivalent magnitude although the month to month variability of the change of heat content is greater than that of the heat exchange. If one assumes that the magnitude of the heat exchange is correct, then a greater decline in heat content than expected from heat exchange tends to occur during fall and winter. Heat content increases, greater than those expected from heat exchange, tend to occur during spring and summer. According to our estimates the heat exchange across the sea surface contributes only 47% to the variance in the rate of change of heat content due to mean temperature change. The remainder must be attributed to advection and diffusion of which the former is probably the dominant process.

Another opportunity to examine the effect of total heat exchange across the sea surface occurred during the time from the fall of 1969 to the spring of 1970 when both the surface temperature and the mean temperature in the layer above σ_t 26 were below the 6-yr average. Cooling began in August, 2 mo earlier than average, and continued until March 1970 (Fig. 18). A greater than average increase in temperature occurred in April 1970 (1.1°C instead of 0.1°C). Throughout this time there is no indication of anomalous heat exchange across the sea surface (Fig. 17). It is interesting to note that in April 1970 there was an increase in the surface salinity that subsequently persisted as a shallow subsurface salinity maximum (Fig. 14). There was no significant change in depth of the thermal structure (σ_t 26) during this month of relatively large surface temperature and salinity changes.

In summary, anomalous heat content changes in the 6-yr series cannot be attributed exclusively to anomalous heat exchange across the sea surface. Possibly, effects of processes >1,000 km upstream in the principal heat loss area of the North Pacific Ocean play an important role in the temperature variability. Unfortunately, heat advection cannot be calculated from the properties that were measured at OWS-V.

The salinity and salt content are important properties in isentropic analysis. Salt budget analyses, however, suffer from the lack of reliable precipitation data. The association of a shallow salinity maximum with the depth of σ_t 26 has already been described. In

addition, there appears to be a seasonal trend in the 6-yr average of the surface salinity with the maximum occurring in late winter and spring and the minimum in late summer and early fall. Departures during individual years are large so that the average of only 6 yr may depart significantly from a long-term mean.

Reed and Elliott (1973) have estimated the precipitation at OWS-V from the present weather code in the standard marine weather reports. Although absolute magnitudes of these estimates may be in doubt, the seasonal trend is probably correct. The mean monthly values are listed in Table 4 together with the estimates of mean evaporation at OWS-V (Husby and Seckel 1975). The evaporation minus precipitation values indicate excess evaporation over precipitation during most of the year with lowest values occurring from April to August when the average salinities are also declining at OWS-V.

As in the case of the surface temperature the effect of advection on the surface salinity may be pronounced.

Table 4.—Estimates of mean evaporation rates (E)¹, mean precipitation rates (P)², and evaporation minus precipitation (E-P), (in millimeters per month) at Ocean Weather Station V.

	J	F	M	A	M	J	J	A	S	O	N	D
E	207	204	152	90	59	51	42	78	128	150	201	213
P	70	62	68	68	59	63	39	30	21	32	64	70
E-P	137	142	84	22	0	-12	3	48	107	118	137	143

¹Husby and Seckel (1975).

²Reed and Elliot (1973).

SUMMARY AND CONCLUSIONS

In this report the oceanographic station data obtained at OWS-V from 1966 to 1971 have been presented in a form that lends itself to an examination of the effects of atmospheric forcing on the water structure. The report complements a previous report on large-scale air-sea interactions at OWS-V (Husby and Seckel 1975). The data processing and analysis procedures have been described. In order to facilitate future analysis of ocean structure, the principal features in the ocean variability have been described and heat budget estimates have been made that point to the important processes affecting the temperature in the upper layer at OWS-V.

Rather than presenting the temperature, salinity, and σ_t as a function of depth, the temperature, salinity, and depth were presented as a function of σ_t which we call the isentropic format. Harmonic analysis was used as a curve-fitting procedure to summarize the isentropic data with a relatively small number of coefficients and to provide smoothed estimates of daily observations. Large gaps in the sampling record in 1966 and 1967 limit the usefulness of the harmonic coefficients during these time periods. Results of the harmonic analyses are presented in the appendix showing the harmonic coefficients for the temperature, salinity, and depth at intervals of 0.2 σ_t .

Presentation of the oceanographic station data in isentropic format shows that the apparently complex




Figure 17.—Heat content change per month (30-day) based on change in mean temperature per month in layer from sea surface to σ_t computed from the harmonic series (see text) (solid line) and heat content change per month contributed by net heat exchange across sea surface (dashed line).




Figure 18.—Change per month (30-day) of the mean temperature from sea surface to σ_t 26.0 at Ocean Weather Station V, computed from the harmonic series (solid line). The 6-yr mean values are indicated by the dotted line.

distribution of properties at OWS-V have some order and permit identification and isolation of different processes that contribute to changes in the distributions. This point is illustrated by the time-series sections of salinity versus σ_t (Fig. 14). At depths greater than σ_t 26 the T-S relationships remain relatively constant with the salinity minimum occurring at a constant σ_t level, σ_t 26.8.

Variations in the depth of sigma- t below σ_t 26 (Fig. 14) reflect changes in the baroclinicity of the area. Deepening of the pycnocline as indicated by the depth of σ_t 26 is associated with an increase in the surface salinity during winter or the appearance of a shallow subsurface salinity maximum at other times of the year. A shallow subsurface salinity maximum is characteristic of the Kuroshio Current. Meridional sections near OWS-V show the pycnocline depth to increase southward. Thus, the associated changes in the salinity of the upper layers above σ_t 26 indicate meandering of the Kuroshio Extension.

The net heat loss across the sea surface at OWS-V during the fall and winter 1967-68 was anomalously high and during 1968-69 anomalously low. These anomalies were not reflected in changes of the heat content in the layer above σ_t 26. In fact, changes in the thickness of the layer are the dominant influence in changes of heat content, obscuring effects of the seasonal variation of the heat exchange across the sea surface. The variation of the heat content of the layer due to a change in the mean temperature (Fig. 17) does reflect the seasonal variation of heat exchange across the sea surface. Again, however, the anomalous heat loss during 1967-68 and 1968-69 are not reflected in anomalous declines of the mean temperature of the upper layer. Anomalously low temperatures beginning in the summer of 1969 and lasting to the spring of 1970 were not caused by anomalous heat exchange. Thus, horizontal advection and possibly, to a lesser extent, diffusion play a dominant role in the temperature variability at OWS-V.

It is interesting to note that the surface divergence as reflected by changes in the depth of σ_t 26 has no annual periodicity. The 6-yr record shows that large baroclinic variability occurs at time scales of more than 55 days with largest variability occurring at the interannual scale.

OWS-V is located in a dynamically active ocean area with the area of highest net annual heat loss in the North Pacific lying over 1,000 km upstream. Relatively large meridional temperature gradients vary seasonally and probably also from month to month. Velocity variations of the Kuroshio Extension at the time scales considered here must also be expected. Consequently, divergence and horizontal advection play an important role in the variability of the observed properties, obscuring the effects of local atmospheric forcing.

The preliminary analyses reported here show that important properties of the ocean structure can be obtained from a monitoring station such as OWS-V. It may be more difficult, however, to relate changes in the structure with local atmospheric forcing. In this case the effects of heat exchange across the sea surface are ob-

scured by other processes. Some of these processes could possibly be determined by measurements that have not been made on weather ships. Obviously, anomalous atmospheric forcing must produce anomalous ocean conditions. It appears that these relationships can be established best if fields of properties in ocean and atmosphere are monitored in order that the effect of divergence and advection can be determined, or that area integrals can be obtained.

ACKNOWLEDGMENTS

We thank A. Bakun, Pacific Environmental Group, NMFS; R. L. Elsberry, Naval Postgraduate School; and S. Tabata, Institute of Ocean Sciences, Sidney, B.C., for reviewing the manuscript and making many useful comments.

LITERATURE CITED

- BARKLEY, R. A.
1968. Oceanographic atlas of the Pacific Ocean. Univ. Hawaii Press, Honolulu, 20 p., 156 figs.
- BERNSTEIN, R. L., and W. B. WHITE.
1977. Zonal variability in the distribution of eddy energy in the mid-latitude North Pacific Ocean. *J. Phys. Oceanogr.* 7:123-126.
- FLEMING, R. H.
1955. Review of the oceanography of the northern Pacific. *Int. North Pac. Fish. Comm., Bull.* 2, 43 p.
- HUSBY, D. M.
1968. Oceanographic observations North Pacific Ocean Station VICTOR 34°00'N, 164°00'E. December 1964—August 1966. U.S. Coast Guard Oceanogr. Rep. 15, CG 373-15, 141 p.
- HUSBY, D. M., and G. R. SECKEL.
1975. Large-scale air-sea interactions at ocean weather station V, 1951-71. U.S. Dep. Commer., NOAA Tech. Rep. NMFS SSRF-696, 44 p.
- JAPAN HYDROGRAPHIC ASSOCIATION.
1975. Marine environmental atlas, northwestern Pacific Ocean - statistics of the oceanographic elements, 164 p.
- JENKINS, G. M., and D. G. WATTS.
1968. Spectral analysis and its applications. Holden-Day Inc., San Francisco, 525 p.
- KAWAI, H.
1972. Hydrography of the Kuroshio Extension. *In* H. Stommel and K. Yoshida (editors), *Kuroshio: physical aspects of the Japan Current*, p. 235-352. Univ. Wash. Press, Seattle.
- MASUZAWA, J.
1972. Water characteristics of the North Pacific central region. *In* H. Stommel and K. Yoshida (editors), *Kuroshio: physical aspects of the Japan Current*, p. 95-127. Univ. Wash. Press, Seattle.
- MONTGOMERY, R. B.
1938. Circulation in upper layers of southern North Atlantic deduced with use of isentropic analysis. *Mass. Inst. Technol., Pap. Phys. Oceanogr. Meteorol.* 6(2), 55 p.
- MONTGOMERY, R. B., and W. S. WOOSTER.
1954. Thermocline anomaly and the analysis of serial oceanographic data. *Deep-Sea Res.* 2:63-70.
- MUROMTSEV, A. M.
1958. The principal hydrological features of the Pacific Ocean. (*Osnovnye cherty gidrologii Tikhogo Okeana.*) (Translated from the Russian by A. Birron and Z. S. Cole in 1963. Israel Program for Scientific Translations, Jerusalem, 417 p. Available from Natl. Tech. Inf. Serv., Springfield, VA 22151, as TT 63-11065.)
- PARR, A. E.
1938. Isopycnal analysis of current flow by means of identifying properties. *J. Mar. Res.* 1:133-154.

- REED, R. K., and W. P. ELLIOTT.
1973. Precipitation at ocean weather stations in the North Pacific. *J. Geophys. Res.* 78:7087-7091.
- REID, J. L., Jr.
1961. On the geostrophic flow at the surface of the Pacific Ocean with respect to the 1,000-decibar surface. *Tellus* 13:489-502.
1965. Intermediate waters of the Pacific Ocean. *John Hopkins Oceanogr. Stud.* 2, 85 p.
- RODEN, G. I.
1972. Temperature and salinity fronts at the boundaries of the sub-arctic-subtropical transition zone in the western Pacific. *J. Geophys. Res.* 77:7175-7187.
- SVERDRUP, H. U., M. W. JOHNSON, and R. H. FLEMING.
1942. *The oceans, their physics, chemistry, and general biology.* Prentice-Hall, N.Y., x + 1087 p.
- TABATA, S.
1965. Variability of oceanographic conditions at ocean station "P" in the northeast Pacific Ocean. *Trans. R. Soc. Can. 3(Ser. 4):*367-418.
- UDA, M.
1938. Researches on "Siome" or current rip in the seas and oceans. *Geophys. Mag.* 11:307-372.
1963. Oceanography of the subarctic Pacific Ocean. *J. Fish. Res. Board Can.* 20:119-179.
- WYRTKI, K.
1975. Fluctuations of the dynamic topography in the Pacific Ocean. *J. Phys. Oceanogr.* 5:450-459.
- YONG, M. Y. Y.
1971. Sea surface temperatures and salinities collected between 1957 and 1969 at nine Pacific monitoring stations. U.S. Dep. Commer., NOAA, NMFS Data Rep. 69, 35 p., on 1 microfiche. [Available Natl. Tech. Int. Serv., 5285 Port Royal Road, Springfield, VA 22151, as COM-72-10099.]

APPENDIX I

Tables of coefficients for the calculation of temperatures, °C, salinities, ‰, and depths, m, at sigma- t levels at Ocean Weather Station V, 1966-71, using the series

$$E(t) = f(1) + M \cdot (t-1) + A_0 + \sum_n \left[A_n \cos \frac{n\omega t}{T} + B_n \sin \frac{n\omega t}{T} \right]$$

$$n = 1, 2, \dots, 12$$

where $f(1)$ = value of property at first day of year,

M = slope of straight line between $f(1)$ and $f(366)$,

$$\omega = \frac{2\pi}{366} \text{ day}^{-1}, \text{ except for 1968 where } \omega = \frac{2\pi}{367} \text{ day}^{-1},$$

t = Julian day of the year

Upper row of Fourier coefficients at each level contains A -coefficients beginning with A_1 . Lower row contains B -coefficients beginning with B_1 .

OMS-V ISENTROPIC SUMMARY, 1966

PROPERTY = TEMP. (C)

FUNDAMENTAL PERIOD = 366 DAYS

LEVEL	F(1)	M	FIRST OBS.	LAST OBS.	A(0)	20.17	5.75	4.82	7.75	12.57	16.07	11.06	6.88	4.24	15.54	1.16	4.47
SURF.	18.01	.0025	4	361	2.17	-196.75	-59.28	-17.68	2.17	20.17	5.75	-11.26	-11.26	-11.26	-11.26	-11.26	-11.26
22.80	25.84	-.0010	229	280	.09	-11.33	88.59	-2.99	1.37	3.37	-1.33	-1.04	1.04	1.04	1.04	1.04	1.04
23.00	24.89	.0003	218	286	.03	-.92	5.06	6.02	-1.86	4.51	-2.01	-1.47	1.47	1.47	1.47	1.47	1.47
23.20	24.41	-.0004	216	286	.08	-7.05	-5.16	2.64	-4.51	1.02	1.91	-1.37	1.37	1.37	1.37	1.37	1.37
23.40	23.97	-.0007	188	287	.07	-8.01	-6.78	2.67	-3.71	1.71	-1.76	-1.47	1.47	1.47	1.47	1.47	1.47
23.60	23.32	-.0007	188	288	.06	-7.16	-3.23	1.20	-1.37	1.37	-1.85	-1.60	1.60	1.60	1.60	1.60	1.60
23.80	22.76	-.0009	186	288	.06	-7.21	-1.09	1.94	-1.42	1.42	-1.70	-1.59	1.59	1.59	1.59	1.59	1.59
24.00	21.97	-.0005	181	288	.06	-6.73	1.74	1.79	-1.32	1.32	-1.53	-1.43	1.43	1.43	1.43	1.43	1.43
24.20	21.12	.0008	179	327	-.03	-1.87	3.33	1.30	-1.58	1.58	-1.39	-1.28	1.28	1.28	1.28	1.28	1.28
24.40	20.66	.0000	176	327	-.01	-3.51	3.65	-4.39	1.46	1.46	-2.55	-2.39	2.39	2.39	2.39	2.39	2.39
24.60	19.86	.0001	170	335	.04	-1.76	1.84	1.88	-1.55	1.55	-1.63	-1.52	1.52	1.52	1.52	1.52	1.52
24.80	18.20	.0020	7	361	.43	-4.03	1.06	-1.84	1.15	1.15	-2.02	-1.94	1.94	1.94	1.94	1.94	1.94
25.00	17.20	.0028	4	361	.46	-4.10	-5.47	-3.57	-2.94	-2.94	-2.37	-2.26	2.26	2.26	2.26	2.26	2.26
25.20	16.28	.0039	4	361	.50	-10.06	6.40	-1.77	2.69	2.69	-2.54	-2.43	2.43	2.43	2.43	2.43	2.43
25.40	15.49	.0034	4	361	.48	-14.87	-9.70	-15.16	-5.38	-5.38	-1.71	-1.62	1.62	1.62	1.62	1.62	1.62
25.60	14.98	.0020	4	361	.17	-13.81	-19.94	-15.54	-4.76	-4.76	4.6	3.52	3.52	3.52	3.52	3.52	3.52
25.80	14.25	.0007	4	361	-.11	-17.32	-14.67	-3.39	-2.18	-2.18	-3.77	-3.68	3.68	3.68	3.68	3.68	3.68
26.00	13.13	-.0002	4	361	-.14	-1.50	-2.83	-4.73	1.19	1.19	-1.65	-1.58	1.58	1.58	1.58	1.58	1.58
26.20	11.72	-.0012	4	361	-.05	-10.45	-1.95	1.82	-2.09	-2.09	-4.06	-3.92	3.92	3.92	3.92	3.92	3.92
26.40	10.00	-.0015	4	361	.05	-13.51	-10.52	1.55	-2.61	-2.61	1.07	1.07	1.07	1.07	1.07	1.07	1.07
26.60	7.68	-.0001	4	361	.16	-12.59	-10.71	-4.95	-2.60	-2.60	-6.74	-6.61	6.61	6.61	6.61	6.61	6.61
26.80	5.90	.0004	4	361	.09	-3.85	-17.09	-2.88	-2.89	-2.89	-4.23	-4.08	4.08	4.08	4.08	4.08	4.08
27.00	4.93	-.0001	4	361	.01	-8.62	-5.29	-3.42	-1.51	-1.51	-3.98	-3.82	3.82	3.82	3.82	3.82	3.82
27.20	3.90	0.0000	4	361	.00	-5.41	-1.87	-5.67	-5.83	-5.83	-4.96	-4.79	4.79	4.79	4.79	4.79	4.79
27.40	3.13	0.0000	4	361	-.07	-10.48	-10.39	-4.84	-1.50	-1.50	-1.57	-1.47	1.47	1.47	1.47	1.47	1.47

OWS-V ISENTROPIC SUMMARY • 1966

PROPERTY = SALINITY

FUNDAMENTAL PERIOD = 366 DAYS

LEVEL	F(1)	M	FIRST OBS.	LAST OBS.	A(0)	FOURIER COEFFICIENTS X 100	FOURIER COEFFICIENTS X 100	FOURIER COEFFICIENTS X 100	FOURIER COEFFICIENTS X 100	FOURIER COEFFICIENTS X 100	FOURIER COEFFICIENTS X 100	FOURIER COEFFICIENTS X 100	FOURIER COEFFICIENTS X 100	FOURIER COEFFICIENTS X 100	FOURIER COEFFICIENTS X 100
27.40	34.74	.0010	4	361	-.01	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
27.20	34.73	.0010	4	361	-.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
27.00	34.70	.0010	4	361	-.01	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
26.80	34.70	.0010	4	361	.02	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
26.60	34.65	.0010	4	361	.04	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
26.40	34.26	.0010	4	361	-.01	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
26.20	34.41	.0010	4	361	-.03	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
26.00	34.50	.0010	4	361	-.04	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
25.80	34.55	.0010	4	361	-.02	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
25.60	34.49	.0010	4	361	.05	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
25.40	34.37	.0010	4	361	.14	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
25.20	34.34	.0010	4	361	.16	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
25.00	34.36	.0010	4	361	.15	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
24.80	34.43	.0010	7	361	.14	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
24.60	34.71	.0010	170	325	.01	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
24.40	34.71	.0010	176	327	-.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
24.20	34.63	.0010	179	327	-.01	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
24.00	34.68	.0010	181	288	.02	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
23.80	34.72	.0010	186	288	.02	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
23.60	34.67	.0010	188	288	.03	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
23.40	34.65	.0010	188	287	.03	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
23.20	34.56	.0010	216	285	.03	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
23.00	34.48	.0010	218	285	.01	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
22.80	34.60	.0010	229	280	.04	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
SURF.	34.42	.0009	4	361	.09	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00

OWS-V ISENTROPIC SUMMARY, 1966

PROPERTY = DEPTH(M.)

FUNDAMENTAL PERIOD = 366 DAYS

FOURIER COEFFICIENTS

LEVEL	F(1)	M	FIRST OBS.	LAST OBS.	A(0)	38	107	13	11	23	12	18	23	05	14
SURF DENSITY	24.83	-0.000	4	361	-0.35	0.33	-0.17	-1.23	-1.18	-0.19	0.16	-1.12	0.23	-0.05	0.14
22.80	2.00	0.603	229	280	-1.94	-5.06	1.86	1.79	-1.59	-0.77	-0.09	0.38	-1.34	-1.15	0.20
23.00	2.00	0.658	218	286	-1.65	-5.60	3.51	-2.19	-1.66	1.05	-1.02	1.00	-1.71	-1.07	1.60
23.20	4.00	0.274	216	286	3.32	-7.98	2.74	-3.05	-1.17	3.05	-1.55	1.22	-1.59	-1.06	0.69
23.40	3.00	0.141	188	287	-3.76	3.47	4.80	-1.55	-1.74	6.43	-1.86	2.02	-1.63	-1.34	0.89
23.60	8.00	0.1370	189	288	-7.89	-6.25	3.65	-2.87	-1.25	-1.32	-1.57	1.50	-1.01	-0.42	0.52
23.80	1.00	0.1699	186	288	-7.62	-1.56	2.14	-2.14	1.44	1.61	-0.87	0.81	-1.26	0.77	0.77
24.00	1.00	0.1835	181	288	-7.78	2.51	-2.86	-2.09	-2.83	1.42	-1.46	1.01	-0.91	1.04	0.88
24.20	5.00	0.1890	179	327	-9.05	1.71	-1.57	-1.30	0.94	1.52	-1.17	1.00	-1.59	1.04	0.77
24.40	21.00	0.1726	176	327	-10.13	1.90	1.44	-1.30	-0.73	1.81	-1.50	1.33	-1.59	1.04	0.89
24.60	20.00	0.1753	170	335	-6.46	3.57	1.85	-1.47	0.07	0.07	-1.49	1.33	-1.36	0.56	0.56
24.80	70.00	0.1342	7	361	-27.21	1.70	1.48	-1.86	-2.99	-0.21	-1.66	1.38	-0.31	0.69	0.69
25.00	35.00	0.3205	4	361	-26.60	4.80	3.33	-2.20	-1.71	0.48	-1.71	1.49	1.08	1.04	0.81
25.20	54.00	0.3781	4	361	-28.44	2.68	3.17	3.53	-2.24	1.77	-0.85	1.58	1.62	0.32	0.32
25.40	70.00	0.4685	4	361	-15.54	2.98	1.77	1.17	-1.63	1.25	-0.87	1.20	3.09	0.73	0.73
25.60	87.00	0.5726	4	361	20.83	2.68	7.51	7.51	-1.68	-1.50	0.34	4.06	2.20	0.97	0.97
25.80	106.00	0.6411	4	361	57.16	8.84	13.33	13.33	-1.37	0.10	1.79	9.06	10.39	2.99	2.99
26.00	134.00	0.6877	4	361	74.26	12.84	28.54	28.54	-1.56	-10.00	17.77	14.73	16.08	5.29	5.29
26.20	177.00	0.7397	4	361	77.82	11.19	29.57	29.57	-1.72	-18.19	17.21	16.81	17.71	-9.31	-9.31
26.40	244.00	0.7260	4	361	77.22	10.94	20.48	20.48	-1.80	-17.92	13.88	13.98	13.86	8.20	8.20
26.60	327.00	0.6712	4	361	79.57	9.70	15.48	15.48	-1.78	-19.95	12.05	12.43	11.76	3.67	3.67
26.80	422.00	0.6712	4	361	73.74	8.44	13.36	13.36	-1.52	-20.95	10.92	10.92	10.91	3.67	3.67
27.00	539.00	0.6630	4	361	73.74	7.39	12.08	12.08	-1.52	-20.95	9.24	9.24	9.24	3.67	3.67
27.20	694.00	0.6356	4	361	78.53	6.15	11.53	11.53	-1.43	-22.51	8.30	8.30	8.30	3.67	3.67
27.40	919.00	0.7507	4	361	88.76	5.04	10.60	10.60	-1.32	-24.43	7.19	7.19	7.19	3.67	3.67

OWS-V ISENTROPIC SUMMARY, 1967

PROPERTY = TEMP. (C)

FUNDAMENTAL PERIOD = 366 DAYS

LEVEL	F(1)	M	FIRST OBS.	LAST OBS.	A(0)	25.01	26.73	18.76	19.85	-10.17	-4.36	FOURIER COEFFICIENTS X 100	12.57	-1.03	14.88	-13.88
SURF.	18.94	.0080	1	328	.66	-2.04	26.73	35.07	2.39	-27.08	-18.36	-7.22	15.43	-1.03	19.19	15.19
22.60	25.84	.0002	268	272	-.01	-.07	22.10	1.06	3.88	-1.37	2.08	-23.18	1.23	-.09	1.10	1.10
22.80	25.21	.0018	268	276	-.17	6.14	-1.25	20.67	3.88	1.14	2.60	1.24	1.86	1.07	1.07	1.07
23.00	25.23	.0001	201	278	-.08	-8.57	19.70	1.19	1.20	-3.89	-1.78	2.05	1.65	1.00	1.00	1.00
23.20	24.40	.0005	197	278	-.08	7.15	-8.32	1.32	1.75	3.89	-2.37	2.76	2.17	-2.00	1.79	1.79
23.40	23.89	-.0002	193	312	-.03	-5.68	6.65	1.77	1.16	-3.15	-1.72	2.47	2.16	-1.51	1.18	1.18
23.60	23.25	.0020	191	313	-.31	12.07	-4.79	1.06	1.07	3.24	-2.82	2.19	2.05	-1.17	1.22	1.22
23.80	21.81	.0025	183	319	-.10	-7.57	1.30	1.30	3.42	-2.78	-.01	2.33	1.71	-1.06	2.27	2.27
24.00	21.05	.0028	183	319	-.11	-16.92	14.35	1.25	3.78	2.08	-5.70	-2.19	1.91	-1.33	2.09	2.09
24.20	20.28	.0036	183	328	-.19	-12.40	11.53	1.30	3.33	2.99	-5.49	-2.06	2.01	-1.50	2.06	2.06
24.40	19.55	.0039	183	328	-.20	-13.98	10.19	1.10	2.86	3.54	-6.07	-4.56	1.40	-2.65	2.37	2.37
24.60	18.99	.0034	183	328	-.16	-4.77	8.32	1.19	3.75	4.46	-5.96	-1.56	1.66	-1.91	2.47	2.47
24.80	18.26	.0030	183	328	-.14	-12.78	8.86	1.25	2.52	4.99	-6.17	-3.59	2.67	-2.00	2.55	2.55
25.00	18.22	.0009	1	328	-.18	-11.24	6.73	1.00	2.69	3.06	-4.78	-1.91	1.18	-1.81	2.25	2.25
25.20	17.69	.0002	1	328	-.18	-8.20	14.08	1.24	2.29	2.19	-3.19	-2.59	1.20	-2.04	2.26	2.26
25.40	16.74	.0003	1	328	-.18	-18.30	17.42	1.30	2.29	5.53	-2.50	-2.59	1.54	-1.65	2.25	2.25
25.60	15.72	-.0004	1	328	-.06	-12.63	13.20	1.20	1.29	5.49	-4.02	-3.32	2.58	-2.46	2.57	2.57
25.80	14.52	-.0010	1	328	-.05	-13.55	10.85	1.05	2.63	1.53	-6.75	-2.96	2.89	-1.89	2.77	2.77
26.00	13.05	-.0006	1	328	.15	-7.53	6.60	1.02	2.91	2.59	-2.08	-2.08	1.19	-1.05	2.48	2.48
26.20	11.31	.0003	1	328	.24	-3.05	5.99	1.05	2.25	2.59	1.54	-1.47	1.22	-1.05	2.05	2.05
26.40	9.60	.0003	1	328	.15	1.00	1.81	1.00	4.15	1.36	-1.67	-4.19	2.02	-1.44	1.80	1.80
26.60	7.65	.0004	1	328	.03	-3.01	5.33	1.03	3.91	3.32	-3.88	-5.69	2.30	-1.11	1.26	1.26
26.80	6.06	.0002	1	328	-.09	-10.07	5.33	1.03	1.43	1.67	-2.78	-2.69	2.05	-4.54	1.32	1.32
27.00	4.80	.0010	1	328	-.19	-5.01	3.25	1.09	3.77	4.04	3.49	-2.54	1.17	-3.17	1.10	1.10
27.20	3.90	.0015	1	328	-.26	1.63	1.62	1.09	2.63	5.22	1.45	-2.53	1.05	-1.06	1.27	1.27
27.40	3.13	.0008	1	328	-.13	7.52	2.15	1.09	1.76	1.20	-2.92	-2.21	1.62	-1.24	1.61	1.61

OWS-V ISENTROPIC SUMMARY • 1967

PROPERTY = SALINITY

FUNDAMENTAL PERIOD = 366 DAYS

LEVEL	F(1)	M	FIRST ORS.	LAST ORS.	A(0)	3.43	2.70	-3.99	2.27	-1.35	3.60	2.22	9	4	7	-1	14	94
SURF.	34.71	.0003	1	328	-.13	3.43	2.70	-3.99	-2.26	-1.35	3.60	2.22	9	4	7	-1	14	94
22.60	34.34	.0001	268	272	-.00	-3.99	-1.62	.15	1.02	-1.15	1.05	.22	2	2	1	-1	1	3
22.40	34.35	.0007	268	276	-.07	3.43	-2.55	2.00	1.55	-1.27	1.05	.26	5	2	1	-1	5	3
23.00	34.62	.0001	201	278	-.03	3.43	3.52	-1.08	1.65	-1.27	1.05	.26	5	2	1	-1	5	3
23.20	34.54	.0002	107	278	-.03	2.41	-2.31	-2.64	2.17	-1.12	1.05	.26	5	2	1	-1	5	3
23.40	34.61	-.0001	193	312	-.01	2.70	-2.35	-2.54	1.62	-1.12	1.05	.26	5	2	1	-1	5	3
23.60	34.54	.0007	191	313	-.11	3.54	-1.15	2.99	1.62	-1.12	1.05	.26	5	2	1	-1	5	3
23.80	34.36	.0000	193	319	-.03	3.83	-3.09	-4.17	1.66	-1.09	1.05	.26	5	2	1	-1	5	3
24.00	34.35	.0010	193	319	-.04	3.56	-2.63	-4.17	1.66	-1.09	1.05	.26	5	2	1	-1	5	3
24.20	34.33	.0013	193	328	-.06	5.04	-4.51	-3.19	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
24.40	34.34	.0013	193	328	-.07	5.18	-5.16	-3.39	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
24.60	34.39	.0012	193	328	-.06	5.32	-4.32	-3.19	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
24.80	34.45	.0010	193	328	-.05	5.61	-3.85	-2.18	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
25.00	34.70	.0003	1	328	-.05	7.96	-2.68	-3.27	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
25.20	34.78	.0011	1	328	-.05	6.07	-1.88	-2.93	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
25.40	34.75	.0001	1	328	-.06	9.38	-5.42	-4.27	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
25.60	34.70	-.0001	1	328	-.02	12.17	-3.99	-1.59	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
25.80	34.60	-.0001	1	328	.00	2.94	-1.02	3.22	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
26.00	34.48	-.0001	1	328	.03	2.64	-1.50	2.24	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
26.20	34.32	.0001	1	328	.05	3.24	-1.32	2.58	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
26.40	34.14	.0002	1	328	.01	1.77	-1.66	1.52	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
26.60	34.05	.0002	1	328	-.01	1.59	-1.55	1.07	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
26.80	34.03	.0001	1	328	-.02	2.02	-1.44	1.07	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
27.00	34.10	.0001	1	328	-.04	1.66	-1.74	1.07	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
27.20	34.22	.0003	1	328	-.05	1.84	-1.85	1.12	1.42	-1.07	1.05	.26	5	2	1	-1	5	3
27.40	34.37	.0001	1	328	-.01	3.90	-1.68	1.10	1.42	-1.07	1.05	.26	5	2	1	-1	5	3

OWS-V ISENTROPIC SUMMARY, 1967
 PROPERTY = DEPTH(M.)

FUNDAMENTAL PERIOD = 365 DAYS

LEVEL	F(1)	M	FIRST ORS.	LAST ORS.	A(0)	02	03	10	05	02	04	FOURIER COEFFICIENTS
24.82	24.82	0.019	1	328	-0.31	0.2	0.0	10	0.5	0.0	0.2	-0.7
26.50	33.00	0.000	268	272	-3.14	3.2	0.7	32	2.9	0.4	0.4	0.5
22.85	35.00	-0.082	268	276	.73	1.5	1.4	14	1.0	1.0	1.0	1.0
23.00	12.00	0.048	201	278	-2.34	3.0	1.8	21	1.0	1.0	1.0	1.0
23.20	14.00	0.713	197	278	-3.92	2.0	1.5	10	1.0	1.0	1.0	1.0
23.40	5.00	1.495	193	312	-8.72	1.7	1.2	8	1.0	1.0	1.0	1.0
27.60	7.00	1.479	191	313	-8.76	1.7	1.3	8	1.0	1.0	1.0	1.0
23.40	1.00	1.459	183	319	-4.73	3.0	1.9	19	1.0	1.0	1.0	1.0
24.00	4.00	1.731	183	319	-8.65	2.0	1.8	12	1.0	1.0	1.0	1.0
24.20	6.00	1.238	187	328	.04	1.9	1.7	12	1.0	1.0	1.0	1.0
24.40	1.00	2.411	187	328	-16.18	1.7	1.6	10	1.0	1.0	1.0	1.0
24.60	15.00	0.263	187	328	-18.73	2.0	1.7	10	1.0	1.0	1.0	1.0
24.80	21.00	0.271	183	328	-19.89	2.1	1.6	10	1.0	1.0	1.0	1.0
25.00	152.00	-0.521	1	328	-46.90	7.4	1.7	10	1.0	1.0	1.0	1.0
25.20	192.00	-0.916	1	328	-60.13	8.7	1.7	10	1.0	1.0	1.0	1.0
26.40	241.00	0.329	1	328	-115.93	13.0	1.6	10	1.0	1.0	1.0	1.0
26.60	296.00	0.822	1	328	-140.95	17.0	1.6	10	1.0	1.0	1.0	1.0
26.80	343.00	0.541	1	328	-139.84	16.0	1.6	10	1.0	1.0	1.0	1.0
26.00	395.00	-0.056	1	328	-129.03	12.0	1.6	10	1.0	1.0	1.0	1.0
26.20	430.00	0.411	1	328	-118.02	12.0	1.6	10	1.0	1.0	1.0	1.0
26.40	492.00	0.217	1	328	-121.98	13.0	1.6	10	1.0	1.0	1.0	1.0
26.60	572.00	0.292	1	328	-136.76	18.0	1.6	10	1.0	1.0	1.0	1.0
26.80	667.00	0.354	1	328	-141.01	15.0	1.6	10	1.0	1.0	1.0	1.0
27.00	791.00	0.457	1	328	-147.75	13.0	1.6	10	1.0	1.0	1.0	1.0
27.20	866.00	0.518	1	328	-139.64	13.0	1.6	10	1.0	1.0	1.0	1.0
27.40	1193.00	0.180	1	328	-97.16	9.0	1.6	10	1.0	1.0	1.0	1.0

OWS-V ISENTROPIC SUMMARY, 1968

PROPERTY = TEMP. (C)

FUNDAMENTAL PERIOD = 367 DAYS

LEVEL	F(I)	M	FIRST OBS.	LAST OBS.	A(O)	15.22	17.21	FOURIER COEFFICIENTS X 100	14.29	1.36	15.17	1.15
SURF.	15.88	.0055	61	362	3.10	-26.49	4.71	-5.26	14.29	1.36	-15.17	-1.15
22.40	26.74	.0012	223	224	-.01	-35.59	-.38	-7.23	-.09	1.20	1.17	-.17
22.60	26.30	-.0014	223	254	.07	2.24	3.35	-.35	1.70	-.17	-1.75	2.46
22.80	26.65	-.0076	203	258	.12	-2.94	1.16	1.01	1.41	-2.07	1.79	-.58
23.00	26.00	-.0045	203	259	.18	-10.31	-1.08	3.73	2.65	-1.60	1.65	2.10
23.20	25.53	-.0077	203	269	.11	-12.65	-1.79	2.75	1.67	-1.59	1.49	1.83
23.40	24.33	-.0013	201	278	-.00	-10.59	2.21	3.24	1.64	1.17	1.05	-.62
23.60	23.29	-.0009	198	302	.06	-4.20	-2.71	1.86	1.58	-1.49	1.20	-.44
23.80	22.28	-.0000	190	302	.07	-6.39	-2.99	-1.84	4.32	-2.49	2.35	-1.77
24.00	21.53	-.0011	189	302	.08	-10.74	-2.89	2.00	5.47	-1.72	1.66	-1.36
24.20	21.41	-.0014	187	302	.11	-11.09	-3.54	-2.00	5.75	-1.80	1.87	-1.82
24.40	23.85	-.0023	170	333	.19	-5.08	3.10	-1.58	4.55	-1.41	2.53	-1.33
24.60	19.97	-.0019	168	334	.21	-3.35	4.29	-1.03	4.51	-1.14	1.51	-1.76
24.80	19.38	-.0021	159	351	.18	-5.52	4.30	-1.05	4.37	-1.20	1.49	-.78
25.00	18.04	-.0011	122	362	.40	-2.88	8.76	-3.11	3.15	-2.02	2.24	-.05
25.20	17.46	-.0015	122	362	.34	-5.01	8.76	-3.07	1.40	-2.08	4.31	2.02
25.40	16.91	-.0020	108	362	.21	-2.91	-2.60	-1.75	2.11	-2.49	3.35	-1.54
25.60	15.88	-.0012	57	362	.12	-6.55	3.74	-5.15	3.74	-3.28	1.96	-1.51
25.80	14.52	-.0011	57	362	.05	-1.55	3.44	-2.16	1.27	-2.00	1.09	-.22
26.00	13.18	-.0004	57	362	-.01	-1.71	3.11	-2.07	1.12	-1.56	1.58	-.15
26.20	11.71	-.0015	57	362	-.02	-1.45	3.26	-2.25	1.28	-1.89	1.47	-.74
26.40	10.05	-.0003	57	362	-.09	-1.91	2.54	-2.32	1.68	-1.57	1.19	1.69
26.60	7.58	-.0011	61	362	.08	-1.55	1.96	-2.37	1.66	-1.39	1.36	1.82
26.80	5.84	-.0004	61	362	.12	-1.05	1.34	-2.42	1.32	-1.34	1.09	3.89
27.00	4.69	-.0002	61	362	.07	-1.17	1.65	-2.44	1.09	-1.69	1.37	1.70
27.20	3.93	-.0012	61	362	-.02	-1.35	1.47	-2.42	1.05	-1.98	1.17	2.36
27.40	3.17	-.0011	61	362	-.05	-1.57	1.16	-2.42	1.04	-1.91	1.07	1.47

OWS-V ISENTROPIC SUMMARY, 1968

PROPERTY = SALINITY

FUNDAMENTAL PERIOD = 367 DAYS

LEVEL	F(1)	M	FIRST OBS.	LAST OBS.	A(0)	3.21	3.86	3.14	2.54	-4.34	FOURIER COEFFICIENTS X 100	1.95	1.97	2.78	2.78	2.96	1.01	1.69	-1.4
SURF.	34.72	-.0005	61	362	.01	-3.21	-3.86	3.14	2.54	-4.34	-1.95	1.95	1.97	-2.50	-2.50	-2.96	1.01	1.69	-1.4
25.40	34.44	.0001	223	224	-.00	-6.60	1.120	2.08	-1.51	1.07	1.97	1.97	.01	.01	.01	.06	.01	.04	-.3
22.60	34.52	-.0006	223	254	.03	2.89	1.08	.16	.17	-.05	.43	.43	.07	.07	.07	.06	.07	.75	1.0
22.80	34.87	-.0015	203	258	.05	6.77	-1.12	2.37	-2.43	1.50	.89	.89	.49	.49	.49	.93	.45	.75	-.1
23.00	34.93	-.0018	203	259	.07	7.79	-3.53	1.59	-1.44	1.69	1.75	1.75	.81	.81	.81	1.39	.69	.45	-.2
23.20	35.00	-.0015	203	269	.04	-2.10	-4.94	3.10	-1.44	1.08	1.81	1.81	1.28	1.28	1.28	1.68	.44	.44	-.4
23.40	34.80	-.0005	201	278	.00	-4.91	-4.55	1.10	-1.69	1.69	1.00	1.00	1.41	1.41	1.41	1.61	.46	.46	-.3
23.60	34.64	-.0003	198	302	.02	-3.21	-1.51	2.60	-1.00	.10	1.13	1.13	.81	.81	.81	1.21	.40	.40	-.2
23.80	34.53	.0000	190	302	.02	-2.77	-2.62	2.93	-1.35	1.35	1.35	1.35	1.28	1.28	1.28	1.48	.58	.58	-.1
24.00	34.51	.0001	199	302	.03	3.71	-4.22	3.53	-1.77	1.83	1.32	1.32	1.81	1.81	1.81	1.58	.61	.61	-.3
24.20	34.74	-.0005	187	302	.04	2.46	-4.35	2.92	-1.65	1.65	1.84	1.84	1.82	1.82	1.82	1.69	.82	.82	-.6
24.40	34.80	-.0008	170	330	.07	2.15	-2.24	1.20	-1.30	.27	1.25	1.25	1.57	1.57	1.57	1.98	.44	.44	-.1
24.60	34.75	-.0007	168	334	.07	1.34	-1.35	1.18	-1.11	.25	.79	.79	1.47	1.47	1.47	1.36	.45	.45	-.4
24.80	34.82	-.0007	159	351	.06	1.80	-2.19	1.26	-1.31	.30	1.05	1.05	1.52	1.52	1.52	1.40	.85	.85	-.4
25.00	34.64	-.0004	122	362	.13	3.52	-1.97	1.34	-1.66	.95	1.90	1.90	1.79	1.79	1.79	1.53	.79	.79	-.1
25.20	34.71	-.0004	122	362	.10	3.19	-1.33	2.05	-1.86	.86	1.66	1.66	1.47	1.47	1.47	1.27	.89	.89	-.1
25.40	34.83	-.0006	138	362	.06	1.33	-1.62	1.28	-1.62	.95	1.66	1.66	1.62	1.62	1.62	1.40	.89	.89	-.1
25.60	34.75	-.0004	57	362	.03	-1.06	-1.92	1.42	-1.55	1.11	1.98	1.98	1.98	1.98	1.98	1.66	.85	.85	-.1
25.80	34.61	0.0000	57	362	.02	-1.14	-1.91	1.13	-1.57	1.41	1.62	1.62	1.62	1.62	1.62	1.40	.85	.85	-.1
26.00	34.52	.0001	57	362	-.01	-1.87	-1.81	1.43	-1.60	1.23	1.62	1.62	1.62	1.62	1.62	1.40	.85	.85	-.1
26.20	34.63	.0001	57	362	-.00	-1.64	-1.42	1.22	-1.95	1.40	1.62	1.62	1.62	1.62	1.62	1.40	.85	.85	-.1
26.40	34.27	.0001	57	362	-.02	-2.28	-2.08	1.48	-2.55	1.79	1.62	1.62	1.62	1.62	1.62	1.40	.85	.85	-.1
26.60	34.04	.0002	61	362	.01	-2.05	-1.49	1.34	-2.58	1.62	1.62	1.62	1.62	1.62	1.62	1.40	.85	.85	-.1
26.80	33.99	.0001	61	362	.02	-1.61	-1.52	1.49	-2.11	1.79	1.62	1.62	1.62	1.62	1.62	1.40	.85	.85	-.1
27.00	34.07	.0001	61	362	.01	-1.49	-1.35	1.76	-2.36	1.49	1.62	1.62	1.62	1.62	1.62	1.40	.85	.85	-.1
27.20	34.23	.0000	61	362	-.01	-1.34	-1.64	1.69	-2.47	1.47	1.62	1.62	1.62	1.62	1.62	1.40	.85	.85	-.1
27.40	34.39	.0000	61	362	-.01	-1.20	-1.05	1.18	-2.43	1.47	1.62	1.62	1.62	1.62	1.62	1.40	.85	.85	-.1

OWS-V ISENTROPIC SUMMARY, 1968

PROPERTY = DEPTH(M.)

FUNDAMENTAL PERIOD = 367 DAYS

FOURIER COEFFICIENTS

LEVEL	F(1)	M	FIRST OBS.	LAST OBS.	A(0)	0.5	00	06	12	18	24	30	36	42	48	54	60	66	72	78	84	90
25.57		-.0017	362	362	-.77	.13	.00	-.06	.01	.03	.00	.05	.00	.07	.00	.05	.00	.04	.00	.01	.03	.00
22.60	6.00	.0191	223	254	-1.00	.12	-.02	-.12	.02	-.02	.02	-.12	.02	-.02	.02	-.12	.02	-.02	.02	-.02	.02	-.02
22.80	1.00	.1230	203	258	-7.48	.11	-.51	-1.05	.24	-.14	.11	-.51	.24	-.14	.11	-.51	.24	-.14	.11	-.51	.24	-.14
23.00	9.00	.0574	203	259	-3.05	.11	-.64	-1.05	.24	-.14	.11	-.51	.24	-.14	.11	-.51	.24	-.14	.11	-.51	.24	-.14
23.20	18.00	0.0000	203	269	-1.62	.12	-.66	-1.05	.24	-.14	.12	-.66	.24	-.14	.12	-.66	.24	-.14	.12	-.66	.24	-.14
23.40	2.00	.0628	201	273	-.39	.14	-.65	-1.05	.24	-.14	.14	-.65	.24	-.14	.14	-.65	.24	-.14	.14	-.65	.24	-.14
23.60	4.00	.0792	199	302	-1.21	.12	-.66	-1.05	.24	-.14	.12	-.66	.24	-.14	.12	-.66	.24	-.14	.12	-.66	.24	-.14
23.80	23.00	.0413	190	302	-1.15	.13	-.66	-1.05	.24	-.14	.13	-.66	.24	-.14	.13	-.66	.24	-.14	.13	-.66	.24	-.14
24.00	13.00	.0847	189	302	-1.73	.13	-.66	-1.05	.24	-.14	.13	-.66	.24	-.14	.13	-.66	.24	-.14	.13	-.66	.24	-.14
24.20	37.00	.0328	187	302	-1.73	.12	-.66	-1.05	.24	-.14	.12	-.66	.24	-.14	.12	-.66	.24	-.14	.12	-.66	.24	-.14
24.40	1.00	.0984	170	330	5.74	.11	-.66	-1.05	.24	-.14	.11	-.66	.24	-.14	.11	-.66	.24	-.14	.11	-.66	.24	-.14
24.60	19.00	.0847	168	334	2.72	.13	-.66	-1.05	.24	-.14	.13	-.66	.24	-.14	.13	-.66	.24	-.14	.13	-.66	.24	-.14
24.80	24.00	.1148	159	351	-.99	.13	-.66	-1.05	.24	-.14	.13	-.66	.24	-.14	.13	-.66	.24	-.14	.13	-.66	.24	-.14
25.00	8.00	.1339	122	362	10.97	.14	-.66	-1.05	.24	-.14	.14	-.66	.24	-.14	.14	-.66	.24	-.14	.14	-.66	.24	-.14
25.20	15.00	.1967	122	362	4.87	.15	-.66	-1.05	.24	-.14	.15	-.66	.24	-.14	.15	-.66	.24	-.14	.15	-.66	.24	-.14
25.40	76.00	.0656	108	362	2.31	.15	-.66	-1.05	.24	-.14	.15	-.66	.24	-.14	.15	-.66	.24	-.14	.15	-.66	.24	-.14
25.60	85.00	.0628	57	362	34.60	.16	-.66	-1.05	.24	-.14	.16	-.66	.24	-.14	.16	-.66	.24	-.14	.16	-.66	.24	-.14
25.80	246.00	-.3373	57	362	15.98	.16	-.66	-1.05	.24	-.14	.16	-.66	.24	-.14	.16	-.66	.24	-.14	.16	-.66	.24	-.14
26.00	350.00	-.5109	57	362	11.01	.17	-.66	-1.05	.24	-.14	.17	-.66	.24	-.14	.17	-.66	.24	-.14	.17	-.66	.24	-.14
26.20	414.00	-.4891	57	362	9.66	.17	-.66	-1.05	.24	-.14	.17	-.66	.24	-.14	.17	-.66	.24	-.14	.17	-.66	.24	-.14
26.40	459.00	-.3934	57	362	16.25	.17	-.66	-1.05	.24	-.14	.17	-.66	.24	-.14	.17	-.66	.24	-.14	.17	-.66	.24	-.14
26.60	470.00	-.1858	51	362	37.74	.17	-.66	-1.05	.24	-.14	.17	-.66	.24	-.14	.17	-.66	.24	-.14	.17	-.66	.24	-.14
26.80	556.00	-.1612	61	362	36.38	.18	-.66	-1.05	.24	-.14	.18	-.66	.24	-.14	.18	-.66	.24	-.14	.18	-.66	.24	-.14
27.00	679.00	-.1913	61	362	37.38	.18	-.66	-1.05	.24	-.14	.18	-.66	.24	-.14	.18	-.66	.24	-.14	.18	-.66	.24	-.14
27.20	872.00	-.2705	61	362	26.42	.18	-.66	-1.05	.24	-.14	.18	-.66	.24	-.14	.18	-.66	.24	-.14	.18	-.66	.24	-.14
27.40	1129.00	-.2377	61	362	20.68	.18	-.66	-1.05	.24	-.14	.18	-.66	.24	-.14	.18	-.66	.24	-.14	.18	-.66	.24	-.14

OWS-V ISENTROPIC SUMMARY, 1969

PROPERTY = TEMP. (C)

FUNDAMENTAL PERIOD = 366 DAYS

LEVEL	F(1)	M	FIRST ORS.	LAST ORS.	A(C)	23.48	23.49	28.43	-22.73	FOURIER COEFFICIENTS X 100	1.78	32.85	94	95
SURF.	17.88	-.0033	3	365	2.82	33.50	33.50	28.43	-22.73	5.04	-3.85	1.78	94	95
22.60	25.19	.0019	209	246	-.03	6.98	6.98	28.43	-18.08	14.04	-3.50	32.85	94	95
22.80	25.33	.0001	190	257	.04	4.99	4.99	28.43	-8.92	-3.47	3.66	1.99	94	95
23.00	24.79	.0008	188	258	-.04	5.47	5.47	28.43	-5.58	5.24	3.89	2.05	94	95
23.20	24.66	-.0019	186	274	.07	4.82	4.82	28.43	-10.42	-3.70	5.13	2.84	94	95
23.40	24.00	-.0021	186	278	.11	7.70	7.70	28.43	-17.15	-3.23	3.84	3.59	94	95
23.60	23.02	-.0012	185	278	.12	11.73	11.73	28.43	-17.50	-2.33	3.44	3.69	94	95
23.80	22.24	-.0010	165	284	.12	5.61	5.61	28.43	-14.57	-2.33	1.97	2.73	94	95
24.00	21.52	-.0005	160	304	.14	13.39	13.39	28.43	-12.03	-1.36	2.95	2.73	94	95
24.20	20.76	-.0002	148	304	.11	19.08	19.08	28.43	-12.99	-1.68	2.07	2.73	94	95
24.40	20.68	-.0023	145	313	.11	17.10	17.10	28.43	-13.06	-1.57	2.83	2.73	94	95
24.60	20.10	-.0024	145	333	.07	22.10	22.10	28.43	-15.07	-1.09	2.57	2.07	94	95
24.80	19.13	-.0019	131	350	.13	15.84	15.84	28.43	-17.55	-1.06	1.94	2.07	94	95
25.00	17.62	.0001	3	357	.45	23.74	23.74	28.43	-12.33	-1.06	3.31	2.07	94	95
25.20	16.97	.0004	3	365	.34	27.04	27.04	28.43	-12.33	-1.06	2.78	2.07	94	95
25.40	16.18	-.0003	3	365	.33	15.75	15.75	28.43	-19.20	-1.06	2.78	2.07	94	95
25.60	15.43	-.0004	3	365	.14	16.75	16.75	28.43	-10.14	-1.06	3.28	2.07	94	95
25.80	14.50	-.0006	3	365	.00	7.83	7.83	28.43	-5.55	-1.06	3.11	2.07	94	95
26.00	13.32	-.0004	3	365	-.04	8.47	8.47	28.43	-4.22	-1.06	3.01	2.07	94	95
26.20	11.89	-.0002	3	365	-.02	2.17	2.17	28.43	3.79	-1.06	2.60	2.07	94	95
26.40	10.15	-.0004	3	365	-.02	4.82	4.82	28.43	-4.47	-1.06	2.44	2.07	94	95
26.60	7.95	-.0006	3	365	.09	7.22	7.22	28.43	-3.23	-1.06	2.57	2.07	94	95
26.80	5.98	-.0002	3	365	.08	3.14	3.14	28.43	-5.09	-1.06	2.76	2.07	94	95
27.00	4.76	-.0002	3	365	.04	2.05	2.05	28.43	-2.55	-1.06	2.74	2.07	94	95
27.20	3.99	-.0007	3	365	.05	1.44	1.44	28.43	-1.88	-1.06	2.19	2.07	94	95
27.40	3.19	-.0003	3	365	-.01	1.27	1.27	28.43	-1.60	-1.06	1.94	2.07	94	95

OMS-V ISENTROPIC SUMMARY • 1969
PROPERTY = SALINITY

FUNDAMENTAL PERIOD = 366 DAYS

LEVEL SURF.	F(1)	M	FIRST OBS.	LAST OBS.	A(0)	FOURIER COEFFICIENTS X 100									
22.60	34.53	-.0000	3	365	.00	2.44	2.65	1.97	-4.12	3.88	-2.94	2.26	-1.77	79	
22.60	34.05	.0008	209	246	-.01	14.71	1.27	1.09	-1.16	1.50	-1.83	1.29	-1.77	69	
22.80	34.39	0.0000	190	257	.02	-9.01	2.07	-2.38	2.22	-1.50	1.00	1.20	-1.19	89	
23.00	34.45	.0003	188	258	-.02	-1.53	1.90	-2.63	-1.75	2.19	-1.72	1.51	-1.13	09	
23.20	34.65	-.0008	186	274	.03	-3.15	1.83	-3.12	-2.13	-1.20	1.57	1.99	-2.16	20	
23.40	34.65	-.0008	186	278	.05	-4.95	1.56	-4.05	-2.14	1.23	-1.44	1.44	-1.33	15	
23.60	34.54	-.0005	185	278	.05	-6.71	1.43	-4.89	-2.11	.87	-1.54	1.06	-1.33	22	
23.80	34.51	-.0004	165	284	.05	-8.14	1.29	-5.63	-2.07	.37	-1.59	.78	-1.42	11	
24.00	34.51	-.0003	160	304	.06	-9.71	1.16	-6.49	-1.95	-.87	-1.45	.82	-1.41	36	
24.20	34.51	-.0001	148	304	.04	-11.31	1.05	-7.36	-1.84	-1.89	-.64	.74	-1.43	45	
24.40	34.74	-.0008	145	313	.04	-13.02	.87	-8.25	-1.74	-1.2	-.45	.39	-1.47	66	
24.60	34.77	-.0007	145	333	.03	-14.79	.73	-9.14	-1.63	-.69	-.25	.33	-1.46	85	
24.80	34.75	-.0007	131	350	.04	-16.64	.55	-10.06	-1.51	-.29	-.04	.23	-1.47	65	
25.00	34.50	.0001	3	357	.14	-18.51	.42	-10.91	-1.39	.70	-.25	.33	-1.47	59	
25.20	34.55	.0001	3	365	.10	-20.44	.29	-11.79	-1.26	1.10	-.22	.69	-1.47	22	
25.40	34.59	-.0001	3	365	.10	-22.41	.17	-12.66	-1.15	1.88	-.04	1.10	-1.47	09	
25.60	34.62	-.0001	3	365	.04	-24.41	.09	-13.53	-1.03	2.50	-.02	1.80	-1.47	22	
25.80	34.61	-.0001	3	365	-.00	-26.44	.02	-14.41	-.92	3.18	-.02	2.48	-1.47	09	
26.00	34.55	-.0001	3	365	-.01	-28.49	.00	-15.30	-.81	3.89	-.01	3.16	-1.47	22	
26.20	34.44	-.0000	3	365	-.00	-30.53	.00	-16.21	-.71	4.62	-.01	3.85	-1.47	09	
26.40	34.30	-.0001	3	365	.00	-32.59	.00	-17.13	-.60	5.37	-.01	4.54	-1.47	09	
26.60	34.11	-.0001	3	365	.01	-34.64	.00	-18.07	-.51	6.09	-.01	5.26	-1.47	09	
26.80	34.01	0.0000	3	365	.01	-36.69	.00	-19.01	-.44	6.84	-.01	5.91	-1.47	09	
27.00	34.09	-.0000	3	365	.00	-38.74	.00	-19.97	-.37	7.58	-.01	6.54	-1.47	09	
27.20	34.23	-.0001	3	365	.01	-40.79	.00	-20.91	-.30	8.33	-.01	7.19	-1.47	09	
27.40	34.39	-.0001	3	365	.00	-42.84	.00	-21.86	-.24	9.08	-.01	7.83	-1.47	09	

OWS-V ISENTROPIC SUMMARY, 1969

PROPERTY = DEPTH(M.)

FUNDAMENTAL PERIOD = 366 DAYS

FOURIER COEFFICIENTS

LEVEL	F(1)	M	FIRST ORS.	LAST ORS.	A(0)	0.79	0.8	0.81	0.82	0.83	0.84	0.85	0.86	0.87	0.88	0.89	0.90	0.91	0.92	0.93	0.94	0.95	0.96	0.97	0.98	0.99	1.00					
SURF	24.96	.0007	3	365	-.71	1.13	-.08	-.00	-.00	.03	.05	.05	.06	.06	.06	.06	.06	.06	.06	.06	.06	.06	.06	.06	.06	.06	.06	.06				
DENSITY	8.00	.0110	209	246	-.40	1.72	.10	-.00	-.00	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04	.04			
	5.00	.0110	190	257	.68	-1.01	.06	-.00	-.00	.11	.15	.15	.15	.15	.15	.15	.15	.15	.15	.15	.15	.15	.15	.15	.15	.15	.15	.15	.15			
	2.00	.0274	188	258	.20	-2.49	1.90	-.00	-.00	.03	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40	.40		
	1.00	.0521	186	274	-.93	-3.70	1.34	-.00	-.00	.03	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51	.51		
	6.00	.0685	186	278	-2.68	-5.32	1.59	-.00	-.00	.03	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	.65	
	4.00	.0164	185	278	4.30	-4.81	1.54	-.00	-.00	.03	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	.77	
	11.00	.0685	165	284	-2.29	-8.21	1.49	-.00	-.00	.03	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	.97	
	17.00	.0384	160	304	-.02	-6.87	2.57	-.00	-.00	.03	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	
	25.00	.0411	148	304	-2.47	-3.93	2.92	-.00	-.00	.03	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	1.77	
	13.00	.0438	145	313	4.20	-2.96	1.46	-.00	-.00	.03	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	1.63	
	18.00	.0849	145	333	-.72	-11.65	1.57	-.00	-.00	.03	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	2.02	
	7.00	.0207	131	350	-9.26	-9.26	1.68	-.00	-.00	.03	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	2.73	
	57.00	.1205	3	357	-27.73	-6.05	1.25	-.00	-.00	.03	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	4.15	
	87.00	-.1507	3	365	-7.18	-7.05	1.25	-.00	-.00	.03	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	5.15	
	100.00	-.0311	3	365	-5.44	-7.26	1.25	-.00	-.00	.03	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	6.15	
	108.00	.0932	3	365	17.04	-35.40	1.25	-.00	-.00	.03	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	7.15	
	124.00	.1205	3	365	41.90	-37.02	1.25	-.00	-.00	.03	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15	8.15
	163.00	.1507	3	365	49.10	-39.56	1.25	-.00	-.00	.03	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15	9.15
	235.00	.1589	3	365	43.80	-26.68	1.25	-.00	-.00	.03	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15	10.15
	315.00	.1616	3	365	36.56	-26.26	1.25	-.00	-.00	.03	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15	11.15
	402.00	.1562	3	365	29.18	-27.06	1.25	-.00	-.00	.03	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15	12.15
	497.00	.1507	3	365	27.45	-28.09	1.25	-.00	-.00	.03	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15	13.15
	609.00	.1945	3	365	24.65	-26.09	1.25	-.00	-.00	.03	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15	14.15
	773.00	.1890	3	365	25.55	-29.66	1.25	-.00	-.00	.03	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15	15.15
	1042.00	.1233	3	365	30.19	-17.71	1.25	-.00	-.00	.03	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15	16.15

OWS-V ISENTROPIC SUMMARY, 1971

PROPERTY = TEMP. (C)

FUNDAMENTAL PERIOD = 366 DAYS

LEVEL	F(1)	M	FIRST OBS.	LAST OBS.	A(C)	FOURIER COEFFICIENTS X 100
SURF.	17.60	-.0013	4	364	3.10	16.13
22.40	26.49	.0001	228	236	-.00	-.35
22.60	25.73	.0005	224	236	-.02	-.30
22.80	25.26	.0004	214	268	-.03	-.27
23.00	24.60	.0002	207	268	-.02	-.25
23.20	23.99	.0004	206	286	-.04	-.24
23.40	23.13	.0012	205	291	-.07	-.23
23.60	22.44	.0019	185	316	-.14	-.22
23.80	21.84	.0033	184	321	-.36	-.21
24.00	20.65	.0039	172	325	-.24	-.20
24.20	20.28	.0017	156	331	-.05	-.19
24.40	20.08	.0005	153	331	-.11	-.18
24.60	19.81	.0002	148	331	-.26	-.17
24.80	19.27	.0000	141	331	-.33	-.16
25.00	17.56	.0027	6	331	-.03	-.15
25.20	16.97	.0001	4	364	.24	-.14
25.40	16.20	-.0007	4	364	.40	-.13
25.60	15.36	-.0004	4	364	.24	-.12
25.80	14.41	.0002	4	364	-.03	-.11
26.00	13.30	-.0005	4	364	-.02	-.10
26.20	11.93	-.0005	4	364	.03	-.09
26.40	10.11	-.0004	4	364	.02	-.08
26.60	7.77	.0004	4	364	.07	-.07
26.80	5.74	.0006	4	364	.06	-.06
27.00	4.71	-.0000	4	364	-.01	-.05
27.20	3.89	.0002	4	364	-.03	-.04
27.40	3.04	.0006	4	364	-.03	-.03

OWS-V ISENTROPIC SUMMARY, 1971

PROPERTY = SALINITY

FUNDAMENTAL PERIOD = 366 DAYS

LEVEL	F(I)	M	FIRST OBS.	LAST OBS.	A(0)	7	5	3	1	175	11	3	100	37	197	1	54	1	83
SURF.	34.51	.0001	4	364	.01	10.33	-7.35	-3.57	-1.57	1.75	-1.11	1.73	2.12	-1.97	-1.57	-1	1.54	-1	1.83
22.40	34.32	.0001	228	236	-.00	14.77	-5.53	-3.47	-1.44	-1.72	1.40	-.71	2.12	-1.97	-1.57	-1	1.54	-1	1.86
25.60	34.28	.0002	224	236	-.01	1.54	-1.28	.08	-1.19	-1.41	1.11	-1.02	2.13	-1.98	-1.58	-1	1.54	-1	1.88
25.90	34.37	.0001	214	268	-.01	1.96	-1.46	.44	1.05	-1.58	1.22	-1.01	2.14	-1.99	-1.59	-1	1.54	-1	1.91
23.00	34.36	.0001	207	268	-.01	1.09	-1.73	.33	1.33	-1.50	1.54	-.07	2.15	-1.99	-1.59	-1	1.54	-1	1.93
23.20	34.38	.0002	206	286	-.02	1.39	-.93	.83	1.05	-1.33	1.93	-.10	2.16	-1.99	-1.59	-1	1.54	-1	1.95
23.40	34.32	.0004	205	291	-.03	2.59	-1.44	1.62	1.11	-1.15	1.13	-.10	2.17	-1.99	-1.59	-1	1.54	-1	1.97
23.60	34.32	.0007	185	316	-.05	3.91	-.73	1.62	1.01	-1.04	1.58	-.14	2.18	-1.99	-1.59	-1	1.54	-1	2.01
23.80	34.37	.0012	184	321	-.13	2.97	-.82	.76	1.52	-1.36	1.40	-.16	2.19	-1.99	-1.59	-1	1.54	-1	2.06
24.00	34.20	.0014	172	325	-.08	10.77	-1.02	1.52	1.58	-2.04	1.10	-.17	2.20	-1.99	-1.59	-1	1.54	-1	2.11
24.20	34.34	.0005	156	331	-.01	5.53	-2.23	3.69	1.18	-2.09	2.20	-.17	2.21	-1.99	-1.59	-1	1.54	-1	2.16
24.40	34.53	.0001	153	331	-.03	4.28	-1.76	2.75	1.44	-1.76	3.15	-.15	2.22	-1.99	-1.59	-1	1.54	-1	2.21
24.60	34.69	.0001	148	331	-.09	6.88	-2.31	2.87	3.08	-1.55	3.91	-.14	2.23	-1.99	-1.59	-1	1.54	-1	2.26
24.80	34.78	-.0000	141	331	-.11	9.40	-2.77	3.55	2.05	-1.95	3.63	-.13	2.24	-1.99	-1.59	-1	1.54	-1	2.31
25.00	34.48	.0009	6	331	-.01	11.29	-3.42	3.28	2.43	-2.08	1.99	-.12	2.25	-1.99	-1.59	-1	1.54	-1	2.36
25.20	34.56	.0000	4	364	-.08	3.06	-3.50	1.32	1.39	-1.78	3.77	-.08	2.26	-1.99	-1.59	-1	1.54	-1	2.41
25.40	34.59	-.0002	4	364	.12	14.93	-2.67	1.66	1.74	-3.02	1.16	-.08	2.27	-1.99	-1.59	-1	1.54	-1	2.46
25.60	34.60	-.0001	4	364	.07	2.07	-3.03	1.39	1.68	-3.55	1.15	-.15	2.28	-1.99	-1.59	-1	1.54	-1	2.51
25.80	34.59	.0000	4	364	-.01	4.64	-3.41	1.16	1.56	-3.58	1.33	-.12	2.29	-1.99	-1.59	-1	1.54	-1	2.56
26.00	34.55	-.0001	4	364	-.01	2.03	-3.49	1.14	1.42	-1.43	1.82	-.11	2.30	-1.99	-1.59	-1	1.54	-1	2.61
26.20	34.43	-.0001	4	364	.01	16.7	-.75	.89	1.55	-3.37	2.03	-.11	2.31	-1.99	-1.59	-1	1.54	-1	2.66
26.40	34.29	-.0001	4	364	.01	1.55	-3.55	1.16	1.42	-3.60	1.99	-.08	2.32	-1.99	-1.59	-1	1.54	-1	2.71
26.60	34.08	.0000	4	364	.02	1.59	-3.52	1.57	1.83	-3.66	1.85	-.05	2.33	-1.99	-1.59	-1	1.54	-1	2.76
26.80	33.98	.0001	4	364	.01	1.40	-3.46	1.44	1.31	-3.51	1.86	-.05	2.34	-1.99	-1.59	-1	1.54	-1	2.81
27.00	34.08	-.0000	4	364	.00	2.27	-3.01	1.44	1.09	-1.16	1.22	-.11	2.35	-1.99	-1.59	-1	1.54	-1	2.86
27.20	34.22	.0000	4	364	-.00	1.48	-3.06	1.05	1.09	1.11	1.14	-.09	2.36	-1.99	-1.59	-1	1.54	-1	2.91
27.40	34.37	.0001	4	364	-.00	1.08	-3.17	.87	1.38	-1.56	1.03	-.08	2.37	-1.99	-1.59	-1	1.54	-1	2.96
						1.75	-3.10	.84	1.09	1.11	1.14	-.07	2.38	-1.99	-1.59	-1	1.54	-1	3.01
						1.02	-3.04	.81	1.24	-1.16	1.22	-.09	2.39	-1.99	-1.59	-1	1.54	-1	3.06

OWS-V ISENTROPIC SUMMARY, 1971

PROPERTY = DEPTH(M.)

FUNDAMENTAL PERIOD = 366 DAYS

LEVEL	F(1)	M	FIRST OBS.	LAST OBS.	A(0)	0.5	0.6	0.4	0.8	0.2	0.3	0.1	0.9	0.0	0.4	0.3	0.0	0.2	0.1
SLPF.	25.01	.0004	4	364	-.83	.59	.06	.04	.08	.02	.02	.00	.04	.00	.00	.00	.00	.00	.00
DENSITY	22.40	-.0082	228	236	.47	1.10	-.18	-.02	-.02	-.02	-.02	-.02	-.02	-.02	-.02	-.02	-.02	-.02	-.02
	22.60	.0082	224	236	-.40	.70	.04	.11	.17	.17	.18	.13	.18	.10	.08	.11	.08	.08	.08
	22.80	.0164	214	268	-.31	-.55	-.04	.12	.07	.22	.25	.12	.21	.10	.19	.14	.15	.15	.15
	23.00	.0329	207	268	-.48	1.98	1.86	.17	.25	.25	.25	.14	.21	.10	.19	.14	.15	.15	.15
	23.20	.0548	206	286	-1.92	1.40	1.16	.16	.46	.46	.46	.28	.38	.20	.28	.28	.28	.28	.28
	23.40	.0712	205	291	-3.95	3.09	.32	.67	.63	.63	.62	.44	.58	.30	.44	.44	.44	.44	.44
	23.60	.0877	185	316	-6.46	5.64	.07	.33	.44	.44	.44	.28	.38	.20	.28	.28	.28	.28	.28
	23.80	.0877	184	321	-2.90	7.38	1.54	.03	.27	.27	.27	.19	.25	.10	.19	.19	.19	.19	.19
	24.00	.1479	172	325	-12.05	3.79	1.57	.58	.69	.69	.69	.44	.58	.30	.44	.44	.44	.44	.44
	24.20	.1425	156	331	-10.88	12.87	1.02	.32	.85	.85	.85	.58	.79	.40	.58	.58	.58	.58	.58
	24.40	.1836	153	331	-12.30	2.04	1.02	.32	.85	.85	.85	.58	.79	.40	.58	.58	.58	.58	.58
	24.60	.2219	148	331	-16.94	3.04	1.02	.32	.85	.85	.85	.58	.79	.40	.58	.58	.58	.58	.58
	24.80	.2356	141	331	-17.67	14.48	1.02	.32	.85	.85	.85	.58	.79	.40	.58	.58	.58	.58	.58
	25.00	.3890	6	331	-43.74	12.80	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	25.20	.0685	4	364	-13.41	11.72	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	25.40	1.09.00	4	364	-22.12	37.04	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	25.60	.0548	4	364	-15.17	58.60	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	25.80	.145.00	4	364	2.22	70.93	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	26.00	.205.00	4	364	-22.12	11.72	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	26.20	.290.00	4	364	-15.17	14.36	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	26.40	.365.00	4	364	-5.31	28.04	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	26.60	.469.00	4	364	-103.62	46.89	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	26.80	.583.00	4	364	-110.67	69.19	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	27.00	.691.00	4	364	-97.07	99.16	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	27.20	.857.00	4	364	-96.53	110.95	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77
	27.40	1.165.00	4	364	-110.78	133.62	1.11	.31	1.11	1.11	1.11	.77	.99	.50	.77	.77	.77	.77	.77

FOURIER COEFFICIENTS

APPENDIX II

Expected values of Ocean Weather Station V temperatures, $^{\circ}\text{C} \times 100$, (top panel); salinities, $\text{‰} \times 100$, (middle panel); and depths, m, (bottom panel) at sigma- t levels evaluated at the first day of each month for the years 1966-71.

Values were computed by using the tabulations of Appendix I in Equation (3). Only the first three harmonics in the series were evaluated for 1966 and 1967. All 12 harmonics in the series were evaluated for 1968 through 1971.

22.0									2674	
								2554	2584	2571
23.0							2509	2483	2469	
							2436	2420	2407	
							2376	2356	2347	2377
						2188	2311	2286	2284	2355
24.0						2130	2298	2290	2221	2270
						2056	2099	2094	2089	2139
				1941	1985	2030	2020	2022	2022	2068
				1825	1846	1889	1888	1683		1920
25.0	1887			1791	1776	1808	1809	1686	1642	
	1769	1717		1727	1696	1730	1734	1726	1760	
	1677	1658	1605	1606	1655	1637	1619	1651	1647	1638
	1574	1550	1565	1585	1595	1547	1537	1553	1540	1542
	1452	1436	1442	1456	1458	1438	1434	1435	1431	1431
26.0	1382	1294	1300	1300	1331	1317	1320	1315	1312	1309
	1127	1143	1152	1154	1163	1164	1156	1173	1178	1160
	951	968	981	985	977	983	978	979	982	979
	753	771	762	778	775	773	758	760	754	778
	594	594	601	596	612	601	588	588	586	601
27.0	476	477	479	475	487	479	474	476	479	482
	390	386	391	390	391	395	391	390	400	393
	315	311	309	313	318	310	310	312	303	308
1000 H	360	376	356	336	331	334	331	321	315	326
	1	32	60	91	121	152	182	213	244	274

22.0									3444	
								3452	3434	3435
23.0							3456	3445	3440	
							3455	3448	3443	
							3456	3449	3445	3459
							3459	3448	3447	3474
24.0						3442	3457	3454	3450	3468
						3444	3458	3456	3453	3470
						3444	3459	3458	3455	3473
						3445	3461	3460	3458	3475
				3459	3448	3464	3463	3461	3461	3475
25.0	3471				3445	3452	3465	3465	3463	3476
	3470				3459	3455	3465	3466	3465	3476
	3476	3456			3456	3458	3466	3468	3465	3476
	3476	3470	3469	3474	3475	3464	3458	3467	3466	3464
	3470	3464	3468	3474	3474	3463	3460	3465	3464	3462
	3468	3457	3459	3463	3463	3457	3456	3457	3457	3457
26.0	3447	3446	3447	3447	3455	3451	3452	3451	3451	3450
	3431	3434	3436	3436	3438	3438	3439	3441	3442	3438
	3418	3420	3423	3423	3421	3423	3422	3420	3422	3424
	3403	3407	3409	3406	3407	3406	3404	3404	3404	3406
	3401	3401	3402	3402	3404	3402	3400	3400	3399	3402
27.0	3409	3409	3409	3408	3410	3409	3409	3410	3410	3407
	3422	3423	3413	3423	3422	3423	3422	3422	3424	3422
	3438	3437	3437	3438	3439	3436	3438	3438	3438	3439
1000 H	3420	3425	3430	3433	3433	3433	3434	3435	3435	3426
	1	32	60	91	121	152	182	213	244	274

22.0									0	
								0	23	32
23.0								17	26	34
								19	30	37
								20	32	39
								23	34	41
24.0						0	24	37	43	50
						7	28	38	45	71
						11	31	41	46	73
						15	36	42	49	88
						20	41	45	52	96
						19	25	48	57	104
25.0	145				14	30	57	62	64	110
	192				20	35	68	72	77	138
	254	238			39	43	84	90	90	166
	317	292	204	149	71	50	50	115	116	108
	368	300	223	183	148	83	61	148	158	135
26.0	418	364	302	248	243	130	108	196	196	170
	454	432	391	307	323	208	191	260	243	228
	504	497	458	400	370	298	275	341	318	301
	586	572	534	494	461	381	359	434	407	374
	678	667	626	582	551	491	479	533	508	470
27.0	791	780	743	698	652	617	608	646	601	584
	945	936	911	868	804	783	784	804	748	748
	1211	1187	1167	1107	1073	1058	1056	1076	1025	980
1000 H	2721	2724	2730	2734	2535	2734	2735	2737	2739	2735
	1	32	60	91	121	152	182	213	244	274

NOAA TECHNICAL REPORTS
NMFS CIRCULAR AND SPECIAL SCIENTIFIC REPORT—FISHERIES
GUIDELINES FOR CONTRIBUTORS

CONTENTS OF MANUSCRIPT

First page. Give the title (as concise as possible) of the paper and the author's name, and footnote the author's affiliation, mailing address, and ZIP code.

Contents. Contains the text headings and abbreviated figure legends and table headings. Dots should follow each entry and page numbers should be omitted.

Abstract. Not to exceed one double-spaced page. Footnotes and literature citations do not belong in the abstract.

Text. See also Form of the Manuscript below. Follow the *U.S. Government Printing Office Style Manual*, 1973 edition. Fish names, follow the American Fisheries Society Special Publication No. 6, *A List of Common and Scientific Names of Fishes from the United States and Canada*, third edition, 1970. Use short, brief, informative headings in place of "Materials and Methods."

Text footnotes. Type on a separate sheet from the text. For unpublished or some processed material, give author, year, title of manuscript, number of pages, and where it is filed—agency and its location.

Personal communications. Cite name in text and footnote. Cite in footnote: John J. Jones, Fishery Biologist, Scripps Institution of Oceanography, La Jolla, CA 92037, pers. commun. 21 May 1977.

Figures. Should be self-explanatory, not requiring reference to the text. All figures should be cited consecutively in the text and their placement indicated in the left-hand margin of the manuscript. Photographs and line drawings should be of "professional" quality—clear and balanced, and can be reduced to 6½ inches (40 picas) for page width or to 3 7/8 inches (19 picas) for single-column width, but no more than 9 inches (54 picas) high. Photos should be printed on glossy paper—sharply focussed, good contrast. Label each figure. List, and typed double spaced, each figure legend. DO NOT SEND original figures to the Scientific Editor; NMFS Scientific Publications Office will request these if they are needed.

Tables. Each table should start on a separate page and should be self-explanatory, not requiring reference to the text. Headings should be short but amply descriptive. Use only horizontal rules. Number table footnotes consecutively across the page from left to right in Arabic numerals; and to avoid confusion with powers, place them to the *left* of the numerals. If the original tables are typed in our format and are clean and legible, these tables will be reproduced as they are. In the text all tables should be cited consecutively and their placement indicated in the left-hand margin of the manuscript.

Acknowledgments. Place at the end of text. Give credit only to those who gave exceptional contributions and *not* to those whose contributions are part of their normal duties.

Literature cited. In text as: Smith and Jones (1977) or (Smith and Jones 1977); if more than one author, list according to years (e.g., Smith 1936; Jones et al. 1975; Doe 1977). All papers referred to in the text should be listed alphabetically by the senior author's surname under the heading "Literature Cited"; only the author's surname and initials are required in the author line. The author is responsible for the accuracy of the literature citations. Abbreviations of names of periodicals and serials should conform to *Biological Abstracts List of Serials with Title Abbreviations*. Format, see recent SSRF or Circular.

Abbreviations and symbols. Common ones, such as mm, m, g, ml, mg, °C (for Celsius), 'r, /s, etc., should be used. Abbreviate units of measures only when used with numerals; periods are rarely used in these abbreviations. But periods are used in et al., vs., e.g., i.e., Wash. (WA is used only with ZIP code), etc. Abbreviations are acceptable in tables and figures where there is lack of space.

Measurements. Should be given in metric units. Other equivalent units may be given in parentheses.

FORM OF THE MANUSCRIPT

Original of the manuscript should be typed double-spaced on white bond paper. Triple space above headings. Send good duplicated copies of manuscript rather than carbon copies. The sequence of the material should be:


- FIRST PAGE
- CONTENTS
- ABSTRACT
- TEXT
- LITERATURE CITED
- TEXT FOOTNOTES
- APPENDIX
- TABLES (each table should be numbered with an Arabic numeral and heading provided)
- LIST OF FIGURE LEGENDS (Entire figure legends, including "Figure" before each number)
- FIGURES

ADDITIONAL INFORMATION

Send ribbon copy and two duplicated copies of the manuscript to:

Dr. Jay C. Quast, Scientific Editor
Northwest and Alaska Fisheries Center
Auke Bay Laboratory
National Marine Fisheries Service, NOAA
P.O. Box 155
Auke Bay, AK 99821

Copies. Fifty copies will be supplied to the senior author and 100 to his organization free of charge.



**UNITED STATES
DEPARTMENT OF COMMERCE**

NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS STAFF
1700 WESTLAKE AVE. N., ROOM 336
SEATTLE, WA 98109
OFFICIAL BUSINESS

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF COMMERCE
COM 210

**THIRD CLASS
BULK RATE**



NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS — Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS — Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS — Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS — Reports containing data, observations, instructions, etc. A partial listing includes data serials, prediction and outlook periodicals, technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS — Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS — Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

**ENVIRONMENTAL SCIENCE INFORMATION CENTER (D822)
ENVIRONMENTAL DATA AND INFORMATION SERVICE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
U.S. DEPARTMENT OF COMMERCE**

6009 Executive Boulevard
Rockville, MD 20852